

Scientific and Technical Information Center

Requester's Full Name: K. Weddington Examiner #: 68082 Date: 6-7-04  
 Art Unit: 1614 Phone Number: 301-272-0587 Serial Number: 091988, 285  
 Mail Box and Bldg/Room Location: \_\_\_\_\_ Results Format Preferred (circle): PAPER DISK E-MAIL

If more than one search is submitted, please prioritize searches in order of need. MEJ

\*\*\*\*\*  
 Please provide a detailed statement of the search topic, and describe as specifically as possible the subject matter to be searched. Include the elected species or structures, keywords, synonyms, acronyms, and registry numbers, and combine with the concept or utility of the invention. Define any terms that may have a special meaning. Give examples or relevant citations, authors, etc, if known. Please attach a copy of the cover sheet, pertinent claims, and abstract.

Title of Invention: \_\_\_\_\_

Inventors (please provide full names): \_\_\_\_\_

Earliest Priority Filing Date: \_\_\_\_\_

*\*For Sequence Searches Only\* Please include all pertinent information (parent, child, divisional, or issued patent numbers) along with the appropriate serial number.*

A composition comprising

- 1) acridine derivatives
- 2) camptothecin derivatives

The acridine derivative is selected from  
 G-F120918, XR 9051 or XR 9576

The camptothecin derivatives selected from

topotecan BNP1350 CPT11  
 Gc211 9-aminocamptothecin  
 DX8951f 9-nitrocamptothecin

STAFF USE ONLY

|   | Type of Search        | Vendors and cost where applicable |
|---|-----------------------|-----------------------------------|
| Searcher: <u>P. Pappal</u>              | NA Sequence (#) _____ | STN <u>✓</u>                      |
| Searcher Phone #: <u>2-2557</u>         | AA Sequence (#) _____ | Dialog _____                      |
| Searcher Location: <u>1665</u>          | Structure (#) _____   | Questel/Orbit _____               |
| Date Searcher Picked Up: <u>6/11/04</u> | Bibliographic _____   | Dr. Link _____                    |
| Date Completed: <u>6/15/04</u>          | Litigation _____      | Lexis/Nexis _____                 |
| Searcher Prep & Review Time _____       | Fulltext _____        | Sequence Systems _____            |
| Clerical Prep Time _____                | Patent Family _____   | WWW/Internet _____                |
| Online Time: _____                      | Other _____           | Other (specify) _____             |

=> b reg

FILE 'REGISTRY' ENTERED AT 10:09:36 ON 15 JUN 2004  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2004 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
provided by InfoChem.

STRUCTURE FILE UPDATES: 14 JUN 2004 HIGHEST RN 693217-50-4  
DICTIONARY FILE UPDATES: 14 JUN 2004 HIGHEST RN 693217-50-4

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 6, 2004

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more  
information enter HELP PROP at an arrow prompt in the file or refer  
to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=> d que l13

|     |    |     |               |        |        |  |
|-----|----|-----|---------------|--------|--------|--|
| L1  | 1  | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | ACRIDINE/CN  |
| L2  | 1  | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | GF 120918/CN   |
| L3  | 1  | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | XR 9051/CN   |
| L4  | 1  | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | CAMPTOTHECIN/CN  |
| L5  | 1  | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | TOPOTECAN/CN   |
| L6  | 1  | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | BNP 1350/CN  |
| L7  | 1  | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | CPT 11/CN  |
| L8  | 1  | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | GG 211/CN  |
| L9  | 1  | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | 9-AMINOCAMPTOTHECIN/CN   |
| L10 | 1  | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | 9-NITROCAMPTOTHECIN/CN   |
| L11 | 1  | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | DX 8951F/CN  |
| L12 | 1  | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | XR 9576/CN   |
| L13 | 12 | SEA | FILE=REGISTRY | ABB=ON | PLU=ON | (L1 OR L2 OR L3 OR L4 OR L5<br>OR L6 OR L7 OR L8 OR L9 OR L10 OR L11 OR L12) |

=> d ide l13 1-12

L13 ANSWER 1 OF 12 REGISTRY COPYRIGHT 2004 ACS on STN  
RN 206873-63-4 REGISTRY  
CN 3-Quinolinecarboxamide, N-[2-[[[4-[2-(3,4-dihydro-6,7-dimethoxy-2(1H)-  
isoquinolinyl)ethyl]phenyl]amino]carbonyl]-4,5-dimethoxyphenyl]- (9CI)  
(CA INDEX NAME)

OTHER NAMES:


CN Tariquidar  
CN XR 9576  
FS 3D CONCORD  
MF C38 H38 N4 O6  
CI COM  
SR CA

LC STN Files: ADISINSIGHT, BIOSIS, CA, CAPLUS, EMBASE, IMSDRUGNEWS,  
IMSPATENTS, IMSRESEARCH, MEDLINE, MRCK\*, PHAR, PROUSDDR, SYNTHLINE,  
TOXCENTER, USAN, USPATFULL

(\*File contains numerically searchable property data)

DT.CA CAPLUS document type: Journal; Patent

RL.P Roles from patents: BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)  
 RL.NP Roles from non-patents: BIOL (Biological study); PREP (Preparation); PROC (Process); PRP (Properties); USES (Uses)



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

19 REFERENCES IN FILE CA (1907 TO DATE)  
 19 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L13 ANSWER 2 OF 12 REGISTRY COPYRIGHT 2004 ACS on STN  
 RN 203923-89-1 REGISTRY  
 CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,  
 4-ethyl-4-hydroxy-11-[2-(trimethylsilyl)ethyl]-, (4S)-(9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,  
 4-ethyl-4-hydroxy-11-[2-(trimethylsilyl)ethyl]-, (S)-

OTHER NAMES:

CN **BNP 1350**

CN DB 172

CN Karenitecin

FS STEREOSEARCH

MF C25 H28 N2 O4 Si

SR CA


LC STN Files: ANABSTR, BIOSIS, CA, CAPLUS, CASREACT, IMSDRUGNEWS,  
 IMSRESEARCH, PROUSDDR, SYNTHLINE, TOXCENTER, USPATFULL

DT.CA Caplus document type: Journal; Patent

RL.P Roles from patents: BIOL (Biological study); PREP (Preparation); PRP (Properties); RACT (Reactant or reagent); USES (Uses)

RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological study); USES (Uses)

Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

24 REFERENCES IN FILE CA (1907 TO DATE)  
24 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L13 ANSWER 3 OF 12 REGISTRY COPYRIGHT 2004 ACS on STN

RN 180422-22-4 REGISTRY

CN Benzamide, N-[4-[2-(3,4-dihydro-6,7-dimethoxy-2(1H)-isoquinolinyl)ethyl]phenyl]-3-[(Z)-[(5Z)-4-methyl-3,6-dioxo-5-(phenylmethylene)piperazinylidene]methyl]-, monohydrochloride (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Benzamide, N-[4-[2-(3,4-dihydro-6,7-dimethoxy-2(1H)-isoquinolinyl)ethyl]phenyl]-3-[[4-methyl-3,6-dioxo-5-(phenylmethylene)piperazinylidene]methyl]-, monohydrochloride, (Z,Z)-

OTHER NAMES:

CN XR 9051

FS STEREOSEARCH

MF C39 H38 N4 O5 . Cl H

SR CA

LC STN Files: CA, CAPLUS, PROUSDDR, SYNTHLINE, TOXCENTER, USPAT2, USPATFULL


DT.CA Caplus document type: Journal; Patent

RL.P Roles from patents: BIOL (Biological study); PREP (Preparation); USES (Uses)

RL.NP Roles from non-patents: BIOL (Biological study); PROC (Process); USES (Uses)

Double bond geometry as shown.

PAGE 1-A



● HCl

PAGE 1-B

— Me

13 REFERENCES IN FILE CA (1907 TO DATE)  
 13 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L13 ANSWER 4 OF 12 REGISTRY COPYRIGHT 2004 ACS on STN

RN 169869-90-3 REGISTRY

CN 10H,13H-Benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-10,13-dione,  
 1-amino-9-ethyl-5-fluoro-1,2,3,9,12,15-hexahydro-9-hydroxy-4-methyl-,  
 (1S,9S)-, monomethanesulfonate (salt) (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 10H,13H-Benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-10,13-dione,  
 1-amino-9-ethyl-5-fluoro-1,2,3,9,12,15-hexahydro-9-hydroxy-4-methyl-,  
 (1S-trans)-, monomethanesulfonate (salt)

OTHER NAMES:

CN DX 8951f

CN Exatecan mesylate

FS STEREOSEARCH

DR 251459-30-0

MF C24 H22 F N3 O4 . C H4 O3 S

SR CA

LC STN Files: ADISINSIGHT, ADISNEWS, BIOSIS, CA, CAPLUS, CASREACT, CIN,  
 IMSDRUGNEWS, IMPATENTS, IMSRESEARCH, PROMT, PROUSDDR, SYNTHLINE,  
 TOXCENTER, USPAT2, USPATFULL

DT.CA CAplus document type: Conference; Journal; Patent

Searched by P. Ruppel


RL.P Roles from patents: BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)  
 RLD.P Roles for non-specific derivatives from patents: BIOL (Biological study); PREP (Preparation); USES (Uses)  
 RL.NP Roles from non-patents: BIOL (Biological study); OCCU (Occurrence); PROC (Process); USES (Uses)

CM 1

CRN 171335-80-1

CMF C24 H22 F N3 O4


Absolute stereochemistry.



CM 2

CRN 75-75-2

CMF C H4 O3 S



50 REFERENCES IN FILE CA (1907 TO DATE)

2 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

51 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L13 ANSWER 5 OF 12 REGISTRY COPYRIGHT 2004 ACS on STN

RN 149882-10-0 REGISTRY

CN 11H-1,4-Dioxino[2,3-g]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-9,12(8H,14H)-dione, 8-ethyl-2,3-dihydro-8-hydroxy-15-[(4-methyl-1-piperazinyl)methyl]-, (8S)-(9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 11H-1,4-Dioxino[2,3-g]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-9,12(8H,14H)-dione, 8-ethyl-2,3-dihydro-8-hydroxy-15-[(4-methyl-1-piperazinyl)methyl]-, (S)-

OTHER NAMES:

CN GG 211

CN GI 147211

CN Lurtotecan

CN NX 211


CN OSI 211

FS STEREOSEARCH

Searched by P. Ruppel

MF C28 H30 N4 O6  
 CI COM  
 SR CA  
 LC STN Files: ADISINSIGHT, ADISNEWS, BIOSIS, BIOTECHNO, CA, CAPLUS, CASREACT, CIN, DDFU, DRUGU, EMBASE, IMSDRUGNEWS, IMSRESEARCH, IPA, PROMT, PROUSDDR, SYNTHLINE, TOXCENTER, USAN, USPAT2, USPATFULL  
 DT.CA CAPLUS document type: Journal; Patent  
 RL.P Roles from patents: BIOL (Biological study); PREP (Preparation); PROC (Process); RACT (Reactant or reagent); USES (Uses)  
 RLD.P Roles for non-specific derivatives from patents: BIOL (Biological study); PREP (Preparation); USES (Uses)  
 RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological study); PREP (Preparation); PROC (Process); RACT (Reactant or reagent); USES (Uses)  
 RLD.NP Roles for non-specific derivatives from non-patents: BIOL (Biological study); PRP (Properties)

Absolute stereochemistry. Rotation (+).



**\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\***

60 REFERENCES IN FILE CA (1907 TO DATE)  
 4 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
 60 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L13 ANSWER 6 OF 12 REGISTRY COPYRIGHT 2004 ACS on STN  
 RN 143664-11-3 REGISTRY  
 CN 4-Acridinecarboxamide, N-[4-[2-(3,4-dihydro-6,7-dimethoxy-2(1H)-isoquinolinyl)ethyl]phenyl]-9,10-dihydro-5-methoxy-9-oxo- (9CI) (CA INDEX NAME)

**OTHER NAMES:**


CN Elacridar  
 CN **GF 120918**  
 CN GG 918  
 CN GW 0918  
 FS 3D CONCORD  
 MF C34 H33 N3 O5  
 CI COM  
 SR CA

LC STN Files: ADISINSIGHT, ANABSTR, BIOSIS, BIOTECHNO, CA, CANCERLIT, CAPLUS, CASREACT, DDFU, DRUGU, EMBASE, IMSDRUGNEWS, IMSRESEARCH, IPA, MEDLINE, PHAR, PROMT, PROUSDDR, SYNTHLINE, TOXCENTER, USAN, USPAT2, USPATFULL

Other Sources: WHO

DT.CA CAPLUS document type: Journal; Patent  
 RL.P Roles from patents: BIOL (Biological study); PREP (Preparation); USES (Uses)  
 RLD.P Roles for non-specific derivatives from patents: BIOL (Biological study); USES (Uses)  
 RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological study); PREP (Preparation); PROC (Process); PRP (Properties); USES (Uses)

PAGE 1-A



PAGE 2-A



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

87 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

88 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L13 ANSWER 7 OF 12 REGISTRY COPYRIGHT 2004 ACS on STN

RN 123948-87-8 REGISTRY

CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,  
 10-[(dimethylamino)methyl]-4-ethyl-4,9-dihydroxy-, (4S)-(9CI) (CA INDEX  
 NAME)

OTHER CA INDEX NAMES:

CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,  
 10-[(dimethylamino)methyl]-4-ethyl-4,9-dihydroxy-, (S)-


Searched by P. Ruppel



## OTHER NAMES:

CN 10-Hydroxy-9-[(dimethylamino)methyl]-(20S)-camptothecin  
CN 9-(N,N-Dimethylaminomethyl)-10-hydroxycamptothecin  
CN Hycamptamine  
CN Hycamptin  
CN NSC 609699  
CN SKF 104864  
CN SKF-S 104864  
CN **Topotecan**  
CN Topotecan lactone  
FS STEREOSEARCH  
DR 133242-28-1, 138121-88-7  
MF C23 H23 N3 O5  
CI COM  
SR CA  
LC STN Files: ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, BIOBUSINESS,  
BIOSIS, BIOTECHNO, CA, CANCERLIT, CAPLUS, CASREACT, CBNB, CEN, CHEMCATS,  
CHEMINFORMRX, CIN, DDFU, DIOGENES, DRUGU, EMBASE, IMSDRUGNEWS,  
IMSPATENTS, IMSRESEARCH, IPA, MEDLINE, MRCK\*, PHAR, PROMT, PROUSDDR, PS,  
RTECS\*, SYNTHLINE, TOXCENTER, USAN, USPAT2, USPATFULL  
(\*File contains numerically searchable property data)  
DT.CA CAplus document type: Book; Conference; Dissertation; Journal; Patent  
RL.P Roles from patents: ANST (Analytical study); BIOL (Biological study);  
PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or  
reagent); USES (Uses)  
RLD.P Roles for non-specific derivatives from patents: ANST (Analytical  
study); BIOL (Biological study); PREP (Preparation); PRP (Properties);  
USES (Uses)  
RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological  
study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU  
(Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); USES  
(Uses)  
RLD.NP Roles for non-specific derivatives from non-patents: BIOL (Biological  
study); PROC (Process); PRP (Properties); USES (Uses)

Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

1049 REFERENCES IN FILE CA (1907 TO DATE)  
29 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
1057 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L13 ANSWER 8 OF 12 REGISTRY COPYRIGHT 2004 ACS on STN

Searched by P. Ruppel

RN 100286-90-6 REGISTRY  
CN [1,4'-Bipiperidine]-1'-carboxylic acid, (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl ester, monohydrochloride (9CI) (CA INDEX NAME)  
OTHER CA INDEX NAMES:  
CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline, [1,4'-bipiperidine]-1'-carboxylic acid deriv.  
CN [1,4'-Bipiperidine]-1'-carboxylic acid, 4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl ester, monohydrochloride, (S)-  
OTHER NAMES:  
CN 7-Ethyl-10-[[4-(1-piperidyl)-1-piperidyl]carbonyloxy]camptothecin hydrochloride  
CN Campto  
CN Camptothecin 11  
CN Camptothecin 11 hydrochloride  
CN **CPT 11**  
CN Irinotecan hydrochloride  
CN Topotecin  
CN U 101440E  
FS STEREOSEARCH  
DR 111348-33-5  
MF C33 H38 N4 O6 . Cl H  
SR CA  
LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BEILSTEIN\*, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CANCERLIT, CAPLUS, CBNB, CHEMCATS, CIN, CSCHEM, DDFU, DIOGENES, DRUGU, EMBASE, IMSCOSEARCH, IMSPATENTS, IMSRESEARCH, IPA, MEDLINE, MRCK\*, PHAR, PROMT, PROUSDDR, PS, RTECS\*, SYNTHLINE, TOXCENTER, USPAT2, USPATFULL  
(\*File contains numerically searchable property data)  
DT.CA Caplus document type: Conference; Journal; Patent  
RL.P Roles from patents: ANST (Analytical study); BIOL (Biological study); PREP (Preparation); PROC (Process); PRP (Properties); USES (Uses)  
RLD.P Roles for non-specific derivatives from patents: BIOL (Biological study); USES (Uses)  
RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)  
RLD.NP Roles for non-specific derivatives from non-patents: BIOL (Biological study); USES (Uses)  
CRN (97682-44-5)

Absolute stereochemistry. Rotation (+).

CCOC(=O)N1CCN(CC1)CC2=CC=CC=C2C3=CC=C4N(C(=O)N5C=CC(=C5)C(=O)OC6C(=O)C(=C(C=C6)C(=O)O)C(=O)OCC6)C3=C4

● HCl

9 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

598 REFERENCES IN FILE CAPLUS (1907 TO DATE)

RN 91421-43-1 REGISTRY

CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14 (4H,12H) -dione,  
10-amino-4-ethyl-4-hydroxy-, (4S) - (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14 (4H,12H) -dione,  
10-amino-4-ethyl-4-hydroxy-, (S)-

OTHER NAMES:

CN 9-Amino-20 (S) -camptothecin

CN 9-Aminocamptothecin

CN NSC 603071

FS STEREOSEARCH

MF C20 H17 N3 O4

| CI | COM |
|----|-----|
|----|-----|

LC STN Files: ADISINSIGHT, ADISNEWS, ANABSTR, BEILSTEIN\*, BIOBUSINESS,  
BIOSIS, CA, CAPLUS, CASREACT, CEN, CHEMCATS, CIN, CSCHEM, IMSDRUGNEWS,  
IMSRESEARCH, IPA, MRCK\*, PHAR, PROMT, PROUSDDR, RTECS\*, SYNTHLINE,  
TOXCENTER, USPAT2, USPATFULL

```
(*File contains numerically searchable property data)
```

DT.CA CAplus document type: Conference; Journal; Patent; Report

RL.P Roles from patents: BIOL (Biological study); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)


RLD.P Roles for non-specific derivatives from patents: BIOL (Biological study); PREP (Preparation); USES (Uses)

RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)

Searched by P. Ruppel

RLD.NP Roles for non-specific derivatives from non-patents: BIOL (Biological study); PREP (Preparation); USES (Uses)

Absolute stereochemistry.



**\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\***

257 REFERENCES IN FILE CA (1907 TO DATE)  
16 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
257 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L13 ANSWER 10 OF 12 REGISTRY COPYRIGHT 2004 ACS on STN

RN 91421-42-0 REGISTRY

CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,  
4-ethyl-4-hydroxy-10-nitro-, (4S)- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,  
4-ethyl-4-hydroxy-10-nitro-, (S)-

OTHER NAMES:

CN 9-Nitro-20(S)-camptothecin

CN **9-Nitrocampthecin**

CN RFS 2000

CN Rubitecan

FS STEREOSEARCH

MF C20 H15 N3 O6

CI COM

LC STN Files: ADISINSIGHT, ADISNEWS, ANABSTR, BEILSTEIN\*, BIOBUSINESS,  
BIOSIS, BIOTECHNO, CA, CAPLUS, CASREACT, CHEMCATS, CIN, CSCHM, DDFU,  
DRUGU, EMBASE, IMSDRUGNEWS, IMSPATENTS, IMSRESEARCH, IPA, MRCK\*, PHAR,  
PROMT, PROUSDDR, SYNTHLINE, TOXCENTER, USAN, USPAT2, USPATFULL  
(\*File contains numerically searchable property data)

DT.CA Caplus document type: Conference; Journal; Patent


RL.P Roles from patents: BIOL (Biological study); PREP (Preparation); PROC  
(Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)

RLD.P Roles for non-specific derivatives from patents: BIOL (Biological  
study); PREP (Preparation); USES (Uses)

RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological  
study); FORM (Formation, nonpreparative); PREP (Preparation); PROC  
(Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)

RLD.NP Roles for non-specific derivatives from non-patents: ANST (Analytical  
study); BIOL (Biological study); PREP (Preparation); USES (Uses)

Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

189 REFERENCES IN FILE CA (1907 TO DATE)

11 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

190 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L13 ANSWER 11 OF 12 REGISTRY COPYRIGHT 2004 ACS on STN

RN 7689-03-4 REGISTRY

CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,  
4-ethyl-4-hydroxy-, (4S)-(9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,  
4-ethyl-4-hydroxy-, (S)-

CN Camptothecin (7CI)

OTHER NAMES:

CN (+)-Camptothecin

CN (+)-Camptothecine

CN (S)-Camptothecin

CN 20(S)-Camptothecin

CN 20(S)-Camptothecine

CN **Camptothecin**

CN d-Camptothecin

CN NSC 94600

FS STEREOSEARCH

DR 30628-51-4, 157405-40-8

MF C20 H16 N2 O4

CI COM

LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BEILSTEIN\*, BIOBUSINESS, BIOSIS,  
BIOTECHNO, CA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CBNB, CEN, CHEMCATS,  
CHEMINFORMRX, CIN, CSCHEM, DDFU, DRUGU, EMBASE, IFICDB, IFIPAT, IFIUDB,  
IPA, MEDLINE, MRCK\*, MSDS-OHS, NAPRALERT, PROMT, PS, RTECS\*, SPECINFO,  
SYNTHLINE, TOXCENTER, USPAT2, USPATFULL

(\*File contains numerically searchable property data)

DT.CA Caplus document type: Book; Conference; Dissertation; Journal; Patent;  
Report

RL.P Roles from patents: ANST (Analytical study); BIOL (Biological study);  
CMBI (Combinatorial study); MSC (Miscellaneous); OCCU (Occurrence); PREP  
(Preparation); PROC (Process); PRP (Properties); RACT (Reactant or  
reagent); USES (Uses)


RLD.P Roles for non-specific derivatives from patents: ANST (Analytical  
study); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation);  
PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES  
(Uses)

RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological

study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses); NORL (No role in record)

RLD.NP Roles for non-specific derivatives from non-patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)

Absolute stereochemistry. Rotation (+).



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

2398 REFERENCES IN FILE CA (1907 TO DATE)

342 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

2413 REFERENCES IN FILE CAPLUS (1907 TO DATE)

1 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

L13 ANSWER 12 OF 12 REGISTRY COPYRIGHT 2004 ACS on STN

RN 260-94-6 REGISTRY

CN **Acridine (8CI, 9CI)** -(CA INDEX NAME)

OTHER NAMES:

CN 10-Azaanthracene

CN 2,3-Benzoquinoline

CN 9-Azaanthracene

CN Benzo[b]quinoline

CN Dibenzo[b,e]pyridine

CN NSC 3408

FS 3D CONCORD

MF C13 H9 N

CI COM, RPS

LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, AQUIRE, BEILSTEIN\*, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CAOLD, CAPLUS, CASREACT, CEN, CHEMCATS, CHEMINFORMRX, CHEMLIST, CIN, CSCHM, CSNB, DDFU, DETHERM\*, DIPPR\*, DRUGU, EMBASE, ENCOMPLIT, ENCOMPLIT2, ENCOMPPAT, ENCOMPPAT2, GMELIN\*, HODOC\*, HSDB\*, IFICDB, IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*, MSDS-OHS, NIOSHTIC, PDLCOM\*, PIRA, PROMT, RTECS\*, SPECINFO, TOXCENTER, TULSA, ULIDAT, USPAT2, USPATFULL, VETU, VTB

(\*File contains numerically searchable property data)

Other Sources: EINECS\*\*, NDSL\*\*, TSCA\*\*


(\*\*Enter CHEMLIST File for up-to-date regulatory information)

DT.CA Caplus document type: Book; Conference; Dissertation; Journal; Patent; Report

RL.P Roles from patents: ANST (Analytical study); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties);

Searched by P. Ruppel

RACT (Reactant or reagent); USES (Uses); NORL (No role in record)  
 RLD.P Roles for non-specific derivatives from patents: ANST (Analytical study); BIOL (Biological study); MSC (Miscellaneous); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)  
 RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological study); CMBI (Combinatorial study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses); NORL (No role in record)  
 RLD.NP Roles for non-specific derivatives from non-patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)



**\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\***

4156 REFERENCES IN FILE CA (1907 TO DATE)  
 421 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
 4159 REFERENCES IN FILE CAPLUS (1907 TO DATE)  
 5 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

=> => b zcaplus

FILE 'ZCAPLUS' ENTERED AT 11:02:53 ON 15 JUN 2004  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS is strictly prohibited.

FILE COVERS 1907 - 15 Jun 2004 VOL 140 ISS 25  
 FILE LAST UPDATED: 14 Jun 2004 (20040614/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d que 121

|    |                            |        |                 |
|----|----------------------------|--------|-----------------|
| L1 | 1 SEA FILE=REGISTRY ABB=ON | PLU=ON | ACRIDINE/CN     |
| L2 | 1 SEA FILE=REGISTRY ABB=ON | PLU=ON | GF 120918/CN    |
| L3 | 1 SEA FILE=REGISTRY ABB=ON | PLU=ON | XR 9051/CN      |
| L4 | 1 SEA FILE=REGISTRY ABB=ON | PLU=ON | CAMPTOTHECIN/CN |

```

L5          1 SEA FILE=REGISTRY ABB=ON  PLU=ON  TOPOTECAN/CN
L6          1 SEA FILE=REGISTRY ABB=ON  PLU=ON  BNP 1350/CN
L7          1 SEA FILE=REGISTRY ABB=ON  PLU=ON  CPT 11/CN
L8          1 SEA FILE=REGISTRY ABB=ON  PLU=ON  GG 211/CN
L9          1 SEA FILE=REGISTRY ABB=ON  PLU=ON  9-AMINOCAMPTOTHECIN/CN
L10         1 SEA FILE=REGISTRY ABB=ON  PLU=ON  9-NITROCAMPTOTHECIN/CN
L11         1 SEA FILE=REGISTRY ABB=ON  PLU=ON  DX 8951F/CN
L12         1 SEA FILE=REGISTRY ABB=ON  PLU=ON  XR 9576/CN
L14        1005 SEA FILE=ZCAPLUS ABB=ON  PLU=ON  L1 (L)?DERIVATIVE?/BI
L15         102 SEA FILE=ZCAPLUS ABB=ON  PLU=ON  (L2 OR L3) OR L12
L16        1106 SEA FILE=ZCAPLUS ABB=ON  PLU=ON  L14 OR L15
L17         258 SEA FILE=ZCAPLUS ABB=ON  PLU=ON  L4 (L)?DERIVATIVE?/BI
L18        1877 SEA FILE=ZCAPLUS ABB=ON  PLU=ON  (L5 OR L6 OR L7 OR L8 OR L9
OR L10 OR L11)
L19        2035 SEA FILE=ZCAPLUS ABB=ON  PLU=ON  L17 OR L18
L20         15 SEA FILE=ZCAPLUS ABB=ON  PLU=ON  L16 AND L19
L21         5 SEA FILE=ZCAPLUS ABB=ON  PLU=ON  L20 AND (PY<=1999 OR PRY<=1999
OR AY<=1999)

```

=> b medl

FILE 'MEDLINE' ENTERED AT 11:03:12 ON 15 JUN 2004

FILE LAST UPDATED: 12 JUN 2004 (20040612/UP). FILE COVERS 1951 TO DATE.

On February 29, 2004, the 2004 MeSH terms were loaded. See HELP RLOAD for details. OLDMEDLINE now back to 1951.

MEDLINE thesauri in the /CN, /CT, and /MN fields incorporate the MeSH 2004 vocabulary. See <http://www.nlm.nih.gov/mesh/> and [http://www.nlm.nih.gov/pubs/techbull/nd03/nd03\\_mesh.html](http://www.nlm.nih.gov/pubs/techbull/nd03/nd03_mesh.html) for a description of changes.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d que 139

```

L22 (      348)SEA FILE=MEDLINE ABB=ON  PLU=ON  ACRIDINE?(2A)DERIVATIVE?
L23 (      460)SEA FILE=MEDLINE ABB=ON  PLU=ON  ACRIDINE/CT,CN OR L22
L24 (       57)SEA FILE=MEDLINE ABB=ON  PLU=ON  "GF 120918"/CN
L25 (        2)SEA FILE=MEDLINE ABB=ON  PLU=ON  "XR 9051"/CN
L26 (       11)SEA FILE=MEDLINE ABB=ON  PLU=ON  TARIQUIDAR/CN
L27 (      526)SEA FILE=MEDLINE ABB=ON  PLU=ON  (L23 OR L24 OR L25 OR L26)
L28 (     3517)SEA FILE=MEDLINE ABB=ON  PLU=ON  CAMPTOTHECIN/CN
L29 (        0)SEA FILE=MEDLINE ABB=ON  PLU=ON  L28 (L) AA
L30 (      860)SEA FILE=MEDLINE ABB=ON  PLU=ON  TOPOTECAN/CN
L31 (        6)SEA FILE=MEDLINE ABB=ON  PLU=ON  "BNP 1350"/CN
L32 (     1564)SEA FILE=MEDLINE ABB=ON  PLU=ON  IRINOTECAN/CN
L33 (        3)SEA FILE=MEDLINE ABB=ON  PLU=ON  GG (2A) 211
L34 (       35)SEA FILE=MEDLINE ABB=ON  PLU=ON  "DX 8951"/CN
L35 (       99)SEA FILE=MEDLINE ABB=ON  PLU=ON  9-AMINO-20-CAMPTOTHECIN/CN
L36 (       73)SEA FILE=MEDLINE ABB=ON  PLU=ON  9-NITROCAMPTOTHECIN/CN
L37 (      312)SEA FILE=MEDLINE ABB=ON  PLU=ON  CAMPTOTHECIN? (2A) ?DERIVATIVE?/
BI
L38 (     2613)SEA FILE=MEDLINE ABB=ON  PLU=ON  (L29 OR L30 OR L31 OR L32 OR
L33 OR L34 OR L35 OR L36 OR L37)
L39         5 SEA FILE=MEDLINE ABB=ON  PLU=ON  L27 AND L38

```

=> b embase



FILE 'EMBASE' ENTERED AT 11:03:22 ON 15 JUN 2004  
COPYRIGHT (C) 2004 Elsevier Inc. All rights reserved.

FILE COVERS 1974 TO 10 Jun 2004 (20040610/ED)

EMBASE has been reloaded. Enter HELP RLOAD for details.

This file contains CAS Registry Numbers for easy and accurate  
substance identification.

=> d que 178

```
L40 (      5550)SEA FILE=EMBASE ABB=ON  PLU=ON  ACRIDINE/CN OR ACRIDINE? OR
      ACRIDINE(2A)?DERIVATIVE?/BI
L41 (      148)SEA FILE=EMBASE ABB=ON  PLU=ON  "GF 120918"/CN OR GF(W)120918
L42 (      147)SEA FILE=EMBASE ABB=ON  PLU=ON  ELACRIDAR/CT
L43 (      180)SEA FILE=EMBASE ABB=ON  PLU=ON  L42 OR L41
L44 (         7)SEA FILE=EMBASE ABB=ON  PLU=ON  "XR 9051"/CN
L45 (         7)SEA FILE=EMBASE ABB=ON  PLU=ON  "XR 9051"/CT
L46 (         7)SEA FILE=EMBASE ABB=ON  PLU=ON  XR(W)9051
L47 (         7)SEA FILE=EMBASE ABB=ON  PLU=ON  (L44 OR L45 OR L46)
L48 (        54)SEA FILE=EMBASE ABB=ON  PLU=ON  "XR 9576"/CN
L49 (        64)SEA FILE=EMBASE ABB=ON  PLU=ON  TARIQUIDAR/CT OR "XR 9576"/CT
L50 (        58)SEA FILE=EMBASE ABB=ON  PLU=ON  XR(W)9576
L51 (        64)SEA FILE=EMBASE ABB=ON  PLU=ON  (L48 OR L49 OR L50)
L52 (       5724)SEA FILE=EMBASE ABB=ON  PLU=ON  L40 OR L43 OR L47 OR L51
L53 (       2502)SEA FILE=EMBASE ABB=ON  PLU=ON  CAMPTOTHECIN/CT
L54 (        839)SEA FILE=EMBASE ABB=ON  PLU=ON  CAMPTOTHECIN(2A)?DERIVATIVE?/BI

L55 (       3448)SEA FILE=EMBASE ABB=ON  PLU=ON  CAMPTOTHECIN?
L56 (       3448)SEA FILE=EMBASE ABB=ON  PLU=ON  (L53 OR L54 OR L55)
L57 (       2675)SEA FILE=EMBASE ABB=ON  PLU=ON  TOPOTECAN/CT
L58 (       2734)SEA FILE=EMBASE ABB=ON  PLU=ON  TOPOTECAN? OR TOPOTECAN(2A)?DER
      IVATIVE?/BI
L59 (       2734)SEA FILE=EMBASE ABB=ON  PLU=ON  L57 OR L58
L60 (         21)SEA FILE=EMBASE ABB=ON  PLU=ON  "BNP 1350"/CT
L61 (         20)SEA FILE=EMBASE ABB=ON  PLU=ON  "BNP 1350"/CN
L62 (         21)SEA FILE=EMBASE ABB=ON  PLU=ON  L60 OR L61
L63 (       1355)SEA FILE=EMBASE ABB=ON  PLU=ON  "CPT 11"/CN
L64 (       4130)SEA FILE=EMBASE ABB=ON  PLU=ON  IRINOTECAN/CT
L65 (       4189)SEA FILE=EMBASE ABB=ON  PLU=ON  IRINOTECAN?
L66 (       4189)SEA FILE=EMBASE ABB=ON  PLU=ON  L64 OR L65
L67 (       4189)SEA FILE=EMBASE ABB=ON  PLU=ON  L66 OR L63
L68 (         13)SEA FILE=EMBASE ABB=ON  PLU=ON  "GG 211"/CN
L69 (         84)SEA FILE=EMBASE ABB=ON  PLU=ON  (LURTOTECAN/CT OR "GG 211"/CT
      OR "GI 147211"/CT)
L70 (         14)SEA FILE=EMBASE ABB=ON  PLU=ON  GG(W)211
L71 (         84)SEA FILE=EMBASE ABB=ON  PLU=ON  L69 OR L70
L72 (         65)SEA FILE=EMBASE ABB=ON  PLU=ON  "DX 8951F"/CN
L73 (        272)SEA FILE=EMBASE ABB=ON  PLU=ON  ?AMINOCAMPTOTHECIN?
L74 (         94)SEA FILE=EMBASE ABB=ON  PLU=ON  ?NITROCAMPTOTHECIN?
L75 (       8543)SEA FILE=EMBASE ABB=ON  PLU=ON  L56 OR L59 OR L62 OR L67 OR
      L68 OR L69 OR L71 OR L72 OR L73 OR L74
L76 (        117)SEA FILE=EMBASE ABB=ON  PLU=ON  L52 AND L75
L77 (      14420)SEA FILE=EMBASE ABB=ON  PLU=ON  DRUG BIOAVAILABILITY/CT
L78 (         10)SEA FILE=EMBASE ABB=ON  PLU=ON  L76 AND L77
```

=> b ipa

FILE 'IPA' ENTERED AT 11:03:35 ON 15 JUN 2004  
COPYRIGHT (C) 2004 American Society of Hospital Pharmacists (ASHP)

FILE COVERS 1970 TO 2 JUN 2004 (20040602/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d que 199

L79 ( 11) SEA FILE=IPA ABB=ON PLU=ON (ACRIDINE/CT OR "ACRIDINE  
DERIVATIVES"/CT OR "ACRIDINE DERIVATIVES, SYNTHESIS"/CT)  
L82 3 SEA FILE=IPA ABB=ON PLU=ON GF-120918/CT  
L83 0 SEA FILE=IPA ABB=ON PLU=ON XR-9051/CT  
L84 0 SEA FILE=IPA ABB=ON PLU=ON XR-9576/CT  
L85 46 SEA FILE=IPA ABB=ON PLU=ON TOPOTECAN/CT  
L86 0 SEA FILE=IPA ABB=ON PLU=ON XR-9576  
L87 75 SEA FILE=IPA ABB=ON PLU=ON (TOPOTECAN/CT OR "TOPOTECAN  
HYDROCHLORIDE"/CT OR "TOPOTECAN HYDROCHLORIDE, APPROVALS"/CT  
OR "TOPOTECAN HYDROCHLORIDE, CHROMATOGRAPHY, LIQUID"/CT OR  
"TOPOTECAN HYDROCHLORIDE, CONTAMINATION"/CT OR "TOPOTECAN  
HYDROCHLORIDE, EQUILIBRIUM CONSTANTS"/CT OR "TOPOTECAN  
HYDROCHLORIDE, FORMULATIONS"/CT OR "TOPOTECAN HYDROCHLORIDE,  
INCOMPATIBILITIES"/CT OR "TOPOTECAN HYDROCHLORIDE, INJECTIONS"/  
CT OR "TOPOTECAN HYDROCHLORIDE, LIPOSOMES"/CT OR "TOPOTECAN  
HYDROCHLORIDE, NEOPLASMS"/CT OR "TOPOTECAN HYDROCHLORIDE,  
OVARIAN NEOPLASMS"/CT OR "TOPOTECAN HYDROCHLORIDE, OVERVIEW"/CT  
OR "TOPOTECAN HYDROCHLORIDE, PHARMACOKINETICS"/CT OR "TOPOTECAN  
HYDROCHLORIDE, REVIEW"/CT OR "TOPOTECAN HYDROCHLORIDE,  
TOXICITY"/CT OR "TOPOTECAN, ADVERSE REACTIONS"/CT OR "TOPOTECAN  
, AVAILABILITY"/CT OR "TOPOTECAN, BREAST NEOPLASMS"/CT OR  
"TOPOTECAN, CARCINOMA"/CT OR "TOPOTECAN, CONTAMINATION"/CT OR  
"TOPOTECAN, DOSAGE"/CT OR "TOPOTECAN, DOSAGE SCHEDULES"/CT OR  
"TOPOTECAN, ENDOMETRIAL NEOPLASMS"/CT OR "TOPOTECAN, GERIATRICS  
"/CT OR "TOPOTECAN, INJECTIONS"/CT OR "TOPOTECAN, LEUKEMIA"/CT  
OR "TOPOTECAN, LUNG NEOPLASMS"/CT OR "TOPOTECAN, MECHANISM OF  
ACTION"/CT OR "TOPOTECAN, MENINGEAL NEOPLASMS"/CT OR "TOPOTECAN  
, NEOPLASMS"/CT OR "TOPOTECAN, OVARIAN NEOPLASMS"/CT OR  
"TOPOTECAN, OVERVIEW"/CT OR "TOPOTECAN, PHARMACOECONOMICS"/CT  
OR "TOPOTECAN, PHARMACOKINETICS"/CT OR "TOPOTECAN, REVIEW"/CT  
OR "TOPOTECAN, SALIVA LEVELS"/CT OR "TOPOTECAN, TOXICITY"/CT  
OR "TOPOTECAN, TOXICITY, ENVIRONMENTAL"/CT)  
L88 0 SEA FILE=IPA ABB=ON PLU=ON BNP-1350/CT  
L89 18 SEA FILE=IPA ABB=ON PLU=ON CPT-11/CT OR "IRINOTECAN HYDROCHLO  
RIDE ANHYDROUS"/CT  
L90 0 SEA FILE=IPA ABB=ON PLU=ON GG-211/CT  
L91 9 SEA FILE=IPA ABB=ON PLU=ON 9-AMINOCAMPTOTHECIN/CT  
L92 2 SEA FILE=IPA ABB=ON PLU=ON 9-NITROCAMPTOTHECIN/CT  
L93 0 SEA FILE=IPA ABB=ON PLU=ON DX-8951F/CT  
L94 4 SEA FILE=IPA ABB=ON PLU=ON L79(L) DERIVATIVE?  
L95 7 SEA FILE=IPA ABB=ON PLU=ON L94 OR (L82 OR L83 OR L84)  
L97 14 SEA FILE=IPA ABB=ON PLU=ON CAMPTOTHECIN(2A) DERIVATIVE?  
L98 107 SEA FILE=IPA ABB=ON PLU=ON L97 OR (L85 OR L86 OR L87 OR L88  
OR L89 OR L90 OR L91 OR L92 OR L93)  
L99 1 SEA FILE=IPA ABB=ON PLU=ON L95 AND L98

=> b biosis

FILE 'BIOSIS' ENTERED AT 11:03:49 ON 15 JUN 2004  
COPYRIGHT (C) 2004 BIOLOGICAL ABSTRACTS INC. (R)

FILE COVERS 1969 TO DATE.  
CAS REGISTRY NUMBERS AND CHEMICAL NAMES (CNs) PRESENT

Searched by P. Ruppel

FROM JANUARY 1969 TO DATE.

RECORDS LAST ADDED: 9 June 2004 (20040609/ED)

FILE RELOADED: 19 October 2003.

=> d que l124

L100( 199)SEA FILE=BIOSIS ABB=ON PLU=ON ACRIDINE/CT OR ("ACRIDINE  
DERIVATIVE"/CT OR "ACRIDINE DERIVATIVES"/CT)  
L101( 432)SEA FILE=BIOSIS ABB=ON PLU=ON ACRIDINE(2A)?DERIVATIVE?/BI  
L102( 568)SEA FILE=BIOSIS ABB=ON PLU=ON L100 OR L101  
L103( 11)SEA FILE=BIOSIS ABB=ON PLU=ON "GF 120918"/CT OR GF-120918/CT  
  
L104( 17)SEA FILE=BIOSIS ABB=ON PLU=ON GF(W)120918  
L105( 17)SEA FILE=BIOSIS ABB=ON PLU=ON L103 OR L104  
L106( 1)SEA FILE=BIOSIS ABB=ON PLU=ON XR-9051/CT  
L107( 2)SEA FILE=BIOSIS ABB=ON PLU=ON XR(W)9051  
L108( 2)SEA FILE=BIOSIS ABB=ON PLU=ON L106 OR L107  
L109( 6)SEA FILE=BIOSIS ABB=ON PLU=ON "XR 9576"/CT  
L110( 8)SEA FILE=BIOSIS ABB=ON PLU=ON XR(W)9576  
L111( 8)SEA FILE=BIOSIS ABB=ON PLU=ON L109 OR L110  
L112( 595)SEA FILE=BIOSIS ABB=ON PLU=ON L102 OR L105 OR L108 OR L111  
L113( 1830)SEA FILE=BIOSIS ABB=ON PLU=ON CAMPTOTHECIN/CT OR ("CAMPTOTHEC  
IN ANALOG"/CT OR "CAMPTOTHECIN ANALOG COMPOUNDS"/CT OR  
"CAMPTOTHECIN ANALOG-CARBOXYMETHYL DEXTRAN CONJUGATE"/CT OR  
"CAMPTOTHECIN ANALOGS"/CT OR "CAMPTOTHECIN ANALOGUE"/CT OR  
"CAMPTOTHECIN ANALOGUES"/CT) OR ("CAMPTOTHECIN COMPOSITION"/CT  
OR "CAMPTOTHECIN COMPOUNDS"/CT OR "CAMPTOTHECIN CONJUGATES"/CT  
OR "CAMPTOTHECIN DERIVATIVE"/CT OR "CAMPTOTHECIN DERIVATIVE  
INTERMEDIATES"/CT OR "CAMPTOTHECIN DERIVATIVE-CARBOXYMETHYL  
DEXTRAN CONJUGATE"/CT OR "CAMPTOTHECIN DERIVATIVE-CARBOXYMETHYL  
DEXTRAN CONJUGATE"/CT OR "CAMPTOTHECIN DERIVATIVE-PLATIN  
DERIVATIVE COMPOSITION"/CT OR "CAMPTOTHECIN DERIVATIVE-POLYMERI  
C CONJUGATES"/CT OR "CAMPTOTHECIN DERIVATIVES"/CT OR "CAMPTOTHE  
CIN DERIVATIVES PREPARATION INTERMEDIATE"/CT OR "CAMPTOTHECIN  
DERIVATIVE"/CT OR "CAMPTOTHECIN DRUG"/CT OR "CAMPTOTHECIN  
DRUGS"/CT OR "CAMPTOTHECIN GLUCOCONJUGATE"/CT OR "CAMPTOTHECIN  
GLYCOCONJUGATE"/CT)  
L114( 359)SEA FILE=BIOSIS ABB=ON PLU=ON CAMPTOTHECIN(2A)?DERIVATIVE?/BI  
  
L115( 1993)SEA FILE=BIOSIS ABB=ON PLU=ON L113 OR L114  
L116( 1452)SEA FILE=BIOSIS ABB=ON PLU=ON TOPOTECAN  
L117( 17)SEA FILE=BIOSIS ABB=ON PLU=ON BNP1350 OR BNP(2A)1350  
L118( 1177)SEA FILE=BIOSIS ABB=ON PLU=ON CPT11 OR CPT(2A)11  
L119( 7)SEA FILE=BIOSIS ABB=ON PLU=ON GG211 OR GG(2A)211  
L120( 43)SEA FILE=BIOSIS ABB=ON PLU=ON DX(2A)8951F OR DX8951F  
L121( 160)SEA FILE=BIOSIS ABB=ON PLU=ON ?AMINOCAMPTOTHECIN?  
L122( 102)SEA FILE=BIOSIS ABB=ON PLU=ON ?NITROCAMPTOTHECIN?  
L123( 4348)SEA FILE=BIOSIS ABB=ON PLU=ON (L115 OR L116 OR L117 OR L118  
OR L119 OR L120 OR L121 OR L122)  
L124 5 SEA FILE=BIOSIS ABB=ON PLU=ON L112 AND L123

=> b wpix

FILE 'WPIX' ENTERED AT 11:04:16 ON 15 JUN 2004  
COPYRIGHT (C) 2004 THOMSON DERWENT

FILE LAST UPDATED: 11 JUN 2004 <20040611/UP>  
MOST RECENT DERWENT UPDATE: 200437 <200437/DW>  
DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

Searched by P. Ruppel

>>> FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,  
PLEASE VISIT:  
[http://www.stn-international.de/training\\_center/patents/stn\\_guide.pdf](http://www.stn-international.de/training_center/patents/stn_guide.pdf) <<<

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES, SEE  
<http://thomsonderwent.com/coverage/latestupdates/> <<<

>>> FOR INFORMATION ON ALL DERWENT WORLD PATENTS INDEX USER  
GUIDES, PLEASE VISIT:  
<http://thomsonderwent.com/support/userguides/> <<<

>>> NEW! FAST-ALERTING ACCESS TO NEWLY-PUBLISHED PATENT  
DOCUMENTATION NOW AVAILABLE IN DERWENT WORLD PATENTS INDEX  
FIRST VIEW - FILE WPIFV. FREE CONNECT HOUR UNTIL 1 MAY 2004.  
FOR FURTHER DETAILS: <http://www.thomsonderwent.com/dwpifv> <<<

>>> NEW! IMPROVE YOUR LITIGATION CHECKING AND INFRINGEMENT  
MONITORING WITH LITALERT. FIRST ACCESS TO RECORDS OF IP  
LAWSUITS FILED IN THE 94 US DISTRICT COURTS SINCE 1973.  
FOR FURTHER DETAILS:  
<http://www.thomsonscientific.com/litalert> <<<

>>> THE DISPLAY LAYOUT HAS BEEN CHANGED TO ACCOMMODATE THE  
NEW FORMAT GERMAN PATENT APPLICATION AND PUBLICATION  
NUMBERS. SEE ALSO:  
<http://www.stn-international.de/archive/stnews/news0104.pdf> <<<

=> d que 1153

|       |       |                  |        |        |   |
|-------|-------|------------------|--------|--------|---|
| L125( | 3517) | SEA FILE=MEDLINE | ABB=ON | PLU=ON | CAMPTOTHECIN/CN   |
| L126( | 0)    | SEA FILE=MEDLINE | ABB=ON | PLU=ON | L125(L)AA   |
| L127( | 860)  | SEA FILE=MEDLINE | ABB=ON | PLU=ON | TOPOTECAN/CN  |
| L128( | 6)    | SEA FILE=MEDLINE | ABB=ON | PLU=ON | "BNP 1350"/CN   |
| L129( | 1564) | SEA FILE=MEDLINE | ABB=ON | PLU=ON | IRINOTECAN/CN   |
| L130( | 3)    | SEA FILE=MEDLINE | ABB=ON | PLU=ON | GG(2A)211   |
| L131( | 35)   | SEA FILE=MEDLINE | ABB=ON | PLU=ON | "DX 8951"/CN  |
| L132( | 99)   | SEA FILE=MEDLINE | ABB=ON | PLU=ON | 9-AMINO-20-CAMPTOTHECIN/CN  |
| L133( | 73)   | SEA FILE=MEDLINE | ABB=ON | PLU=ON | 9-NITROCAMPTOTHECIN/CN  |
| L134( | 312)  | SEA FILE=MEDLINE | ABB=ON | PLU=ON | CAMPTOTHECIN?(2A)?DERIVATIVE?/<br>BI                                      |
| L135( | 2613) | SEA FILE=MEDLINE | ABB=ON | PLU=ON | (L126 OR L127 OR L128 OR L129<br>OR L130 OR L131 OR L132 OR L133 OR L134) |
| L136( | 187)  | SEA FILE=WPIX    | ABB=ON | PLU=ON | (ACRIDINE(W)?DERIVATIVE?)/BIX   |
| L137( | 9)    | SEA FILE=WPIX    | ABB=ON | PLU=ON | (GF120918 OR GF(2A)120918)/BIX  |
| L138( | 4)    | SEA FILE=WPIX    | ABB=ON | PLU=ON | (XR9051 OR XR(2A)9051)/BIX  |
| L139( | 4)    | SEA FILE=WPIX    | ABB=ON | PLU=ON | (XR9576 OR XR(2A)9576)/BIX  |
| L140( | 197)  | SEA FILE=WPIX    | ABB=ON | PLU=ON | L136 OR L137 OR L138 OR L139  |
| L141( | 685)  | SEA FILE=WPIX    | ABB=ON | PLU=ON | CAMPTOTHECIN  |
| L142( | 222)  | SEA FILE=WPIX    | ABB=ON | PLU=ON | CAMPTOTHECIN(2A)?DERIVATIVE?/BIX  |
| L143( | 251)  | SEA FILE=WPIX    | ABB=ON | PLU=ON | TOPOTECAN   |
| L144( | 1)    | SEA FILE=WPIX    | ABB=ON | PLU=ON | BNP1350 OR BNP(2A)1350  |
| L145( | 100)  | SEA FILE=WPIX    | ABB=ON | PLU=ON | CPT11 OR CPT(2A)11  |
| L146( | 3)    | SEA FILE=WPIX    | ABB=ON | PLU=ON | GG211 OR GG(2A)211  |
| L147( | 6)    | SEA FILE=WPIX    | ABB=ON | PLU=ON | DX8951F OR DX(2A)8951F  |
| L148( | 31)   | SEA FILE=WPIX    | ABB=ON | PLU=ON | ?AMINOCAMPTOTHECIN?   |
| L149( | 32)   | SEA FILE=WPIX    | ABB=ON | PLU=ON | ?NITROCAMPTOTHECIN?   |
| L150( | 230)  | SEA FILE=WPIX    | ABB=ON | PLU=ON | (L127 OR L128 OR L129 OR L130 OR<br>L131 OR L132 OR L133 OR L134 OR L135) |
| L151( | 865)  | SEA FILE=WPIX    | ABB=ON | PLU=ON | (L141 OR L142 OR L143 OR L144 OR  |

L145 OR L146 OR L147 OR L148 OR L149)  
L152( 871)SEA FILE=WPIX ABB=ON PLU=ON L150 OR L151  
L153 3 SEA FILE=WPIX ABB=ON PLU=ON L140 AND L152

=> dup rem l124 l39 l21 l78 l99 l153  
FILE 'BIOSIS' ENTERED AT 11:05:39 ON 15 JUN 2004  
COPYRIGHT (C) 2004 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'MEDLINE' ENTERED AT 11:05:39 ON 15 JUN 2004

FILE 'ZCAPLUS' ENTERED AT 11:05:39 ON 15 JUN 2004  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'EMBASE' ENTERED AT 11:05:39 ON 15 JUN 2004  
COPYRIGHT (C) 2004 Elsevier Inc. All rights reserved.

FILE 'IPA' ENTERED AT 11:05:39 ON 15 JUN 2004  
COPYRIGHT (C) 2004 American Society of Hospital Pharmacists (ASHP)

FILE 'WPIX' ENTERED AT 11:05:39 ON 15 JUN 2004  
COPYRIGHT (C) 2004 THOMSON DERWENT  
PROCESSING COMPLETED FOR L124  
PROCESSING COMPLETED FOR L39  
PROCESSING COMPLETED FOR L21  
PROCESSING COMPLETED FOR L78  
PROCESSING COMPLETED FOR L99  
PROCESSING COMPLETED FOR L153  
L154 25 DUP REM L124 L39 L21 L78 L99 L153 (4 DUPLICATES REMOVED)

=> d all l154 1-25

L154 ANSWER 1 OF 25 MEDLINE on STN  
AN 2004082800 MEDLINE  
DN PubMed ID: 14973080  
TI Pheophorbide a is a specific probe for ABCG2 function and inhibition.  
AU Robey Robert W; Steadman Kenneth; Polgar Orsolya; Morisaki Kuniaki;  
Blayney Margaret; Mistry Prakash; Bates Susan E  
CS Cancer Therapeutics Branch, Center for Cancer Research, National Cancer  
Institute, National Institutes of Health, Building 10, Rm. 12C203, 9000  
Rockville Pike, Bethesda, MD 20892, USA.. robeyr@mail.nih.gov  
SO Cancer research, (2004 Feb 15) 64 (4) 1242-6.  
Journal code: 2984705R. ISSN: 0008-5472.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200404  
ED Entered STN: 20040220  
Last Updated on STN: 20040403  
Entered Medline: 20040402  
AB Pheophorbide a (PhA), a chlorophyll catabolite, was shown to be an ABCG2  
substrate based on Abcg2(-/-) knockout mouse studies (J. W. Jonker et  
al., Proc. Natl. Acad. Sci. USA, 99: 15649-15654, 2002). We developed  
a functional assay for ABCG2 using PhA and the ABCG2 inhibitor  
fumitremorgin C. In selected cell lines expressing high levels of  
P-glycoprotein, multidrug resistance-associated protein 1, or ABCG2, PhA  
transport was observed only in cells expressing ABCG2. Fumitremorgin

C-inhibitable PhA transport was found to correlate with cell surface ABCG2 expression as measured by the anti-ABCG2 antibody 5D3. We found that 100 micro M of the cyclin-dependent kinase inhibitor UCN-01 or 1 micro M of the P-glycoprotein inhibitor tariquidar inhibited ABCG2-mediated PhA transport. In 4-day cytotoxicity assays, ABCG2-mediated resistance to SN-38 and topotecan was abrogated in ABCG2-transfected HEK-293 cells treated with 1 micro M tariquidar, and ABCG2-transfected cells were 6-7-fold resistant to UCN-01. PhA is an ABCG2-specific substrate with potential value in measuring ABCG2 function and expression in clinical samples.

CT Check Tags: Human; Support, U.S. Gov't, P.H.S.

\*ATP-Binding Cassette Transporters: AI, antagonists & inhibitors

\*ATP-Binding Cassette Transporters: PH, physiology

Breast Neoplasms: DT, drug therapy

\*Camptothecin: AA, analogs & derivatives

Camptothecin: PD, pharmacology

\*Chlorophyll: AA, analogs & derivatives

\*Chlorophyll: ME, metabolism

Drug Resistance, Neoplasm

\*Neoplasm Proteins: AI, antagonists & inhibitors

\*Neoplasm Proteins: PH, physiology

Quinolines: PD, pharmacology

\*Staurosporine: AA, analogs & derivatives

Staurosporine: PD, pharmacology

Topotecan: PD, pharmacology

RN 112953-11-4 (7-hydroxystaurosporine); 123948-87-8 (Topotecan);

1406-65-1 (Chlorophyll); 15664-29-6 (pheophorbide a); 206873-63-4 (tariquidar); 62996-74-1 (Staurosporine); 7689-03-4 (Camptothecin); 86639-52-3 (7-ethyl-10-hydroxycamptothecin)

CN 0 (ABCG2 protein, human); 0 (ATP-Binding Cassette Transporters); 0 (Neoplasm Proteins); 0 (Quinolines)

L154 ANSWER 2 OF 25 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.  
on STN

AN 2004105539 EMBASE

TI Drug transport at the blood-brain barrier and the choroid plexus.

AU Graff C.L.; Pollack G.M.

CS G.M. Pollack, Division of Drug Delivery, Disposition School of Pharmacy, University of North Carolina, Kerr Hall, Chapel Hill, NC 27599-7360, United States. gary\_pollack@unc.edu

SO Current Drug Metabolism, (2004) 5/1 (95-108).

Refs: 170

ISSN: 1389-2002 CODEN: CDMUBU

CY Netherlands

DT Journal; General Review

FS 008 Neurology and Neurosurgery

029 Clinical Biochemistry

030 Pharmacology

037 Drug Literature Index

LA English

SL English

AB The blood-brain barrier (BBB) and blood-CSF barrier (BCSFB) represent the main interfaces between the central nervous system (CNS) and the peripheral circulation. Drug exposure to the CNS is dependent on a variety of factors, including the physical barrier presented by the BBB and the BCSFB and the affinity of the substrate for specific transport systems located at both of these interfaces. It is the aggregate effect of these factors that ultimately determines the total CNS exposure, and thus pharmacological efficacy, of a drug or drug candidate. This review discusses the anatomical and biochemical barriers presented to solute

access to the CNS. In particular, the important role played by various efflux transporters in the overall barrier function is considered in detail, as current literature suggests that efflux transport likely represents a key determinant of overall CNS exposure for many substrates. Finally, it is important to consider not only the net delivery of the agent to the CNS, but also the ability of the agent to access the relevant target site within the CNS. Potential approaches to increasing both net CNS and target-site exposure, when such exposure is dictated by efflux transport, are considered. .COPYRG. 2004 Bentham Science Publishers Ltd.

CT Medical Descriptors:

- \*drug transport
- \*blood brain barrier
- \*choroid plexus
- biochemistry
- solute
- central nervous system function
- drug exposure
- drug delivery system
- drug targeting
- blood cerebrospinal fluid barrier
- drug mechanism
- drug efficacy
- pain: DT, drug therapy
- drug penetration
- dose response
- drug absorption
- drug disposition
- drug clearance
- drug bioavailability**
- drug protein binding
- drug sensitivity
- transport kinetics
- human
- nonhuman
- review

Drug Descriptors:

- amino acid transporter: EC, endogenous compound
- levodopa: PD, pharmacology
- gabapentin: PD, pharmacology
- glucose transporter 1: EC, endogenous compound
- metenkephalin: PD, pharmacology
- glycoprotein P
- enkephalin[2,5 dextro penicillamine]: DV, drug development
- enkephalin[2,5 dextro penicillamine]: DO, drug dose
- enkephalin[2,5 dextro penicillamine]: DT, drug therapy
- enkephalin[2,5 dextro penicillamine]: PK, pharmacokinetics
- enkephalin[2,5 dextro penicillamine]: PD, pharmacology
- enkephalin[2,5 dextro penicillamine]: IV, intravenous drug administration
- elacridar: PD, pharmacology**
- loperamide: CB, drug combination
- loperamide: CR, drug concentration
- loperamide: IT, drug interaction
- loperamide: PK, pharmacokinetics
- loperamide: PD, pharmacology
- quinidine: CB, drug combination
- quinidine: IT, drug interaction
- quinidine: PD, pharmacology
- breast cancer resistance protein: EC, endogenous compound
- topotecan: PK, pharmacokinetics**
- topotecan: PO, oral drug administration**

mitoxantrone: PK, pharmacokinetics  
 RNA directed DNA polymerase inhibitor: PD, pharmacology  
 multidrug resistance protein: EC, endogenous compound  
 probenecid: PD, pharmacology  
 indometacin: PD, pharmacology  
 mercaptopurine: PD, pharmacology  
 tioguanine: PD, pharmacology  
 adefovir: PD, pharmacology  
 organic cation transporter: EC, endogenous compound  
 carnitine: PD, pharmacology  
 mepyramine: PD, pharmacology  
 organic anion transporter: EC, endogenous compound  
 digoxin: PD, pharmacology  
 estrone sulfate: PD, pharmacology  
 azathioprine: PD, pharmacology  
 tricyclic antidepressant agent: PD, pharmacology  
 penicillin G: PD, pharmacology  
 unindexed drug

RN (levodopa) 59-92-7; (gabapentin) 60142-96-3; (glucose transporter 1) 172077-08-6; (metenkephalin) 58569-55-4; (enkephalin[2,5 dextro penicillamine]) 88373-73-3, 88381-29-7; (elacridar) 143664-11-3; (loperamide) 34552-83-5, 53179-11-6; (quinidine) 56-54-2; (topotecan) 119413-54-6, 123948-87-8; (mitoxantrone) 65271-80-9, 70476-82-3; (multidrug resistance protein) 149200-37-3, 208997-77-7; (probenecid) 57-66-9; (indometacin) 53-86-1, 74252-25-8, 7681-54-1; (mercaptopurine) 31441-78-8, 50-44-2, 6112-76-1; (tioguanine) 154-42-7; (adefovir) 106941-25-7; (carnitine) 461-06-3, 541-15-1, 56-99-5; (mepyramine) 6036-95-9, 91-84-9; (digoxin) 20830-75-5, 57285-89-9; (estrone sulfate) 438-67-5, 481-97-0; (azathioprine) 446-86-6; (penicillin G) 1406-05-9, 61-33-6  
 CN **Gf 120918**

L154 ANSWER 3 OF 25 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.  
 on STN

AN 2004028443 EMBASE

TI Multidrug resistance in cancer chemotherapy and xenobiotic protection mediated by the half ATP-binding cassette transporter ABCG2.

AU Han B.; Zhang J.-T.

CS J.-T. Zhang, Dept. of Pharmacology and Toxicology, IUCC, Indiana Univ. School of Medicine, 1044 W. Walnut Street, Indianapolis, IN 46202, United States. jianzhan@iupui.edu

SO Current Medicinal Chemistry - Anti-Cancer Agents, (2004) 4/1 (31-42).  
 Refs: 74

ISSN: 1568-0118 CODEN: CMCACI

CY Netherlands

DT Journal; General Review

FS 016 Cancer

029 Clinical Biochemistry

030 Pharmacology

037 Drug Literature Index

LA English

SL English

AB ABCG2, also termed BCRP/MXR/ABCP, is a half ATP-binding cassette (ABC) transporter expressed on plasma membranes. ABCG2 was independently cloned from placenta as well as cell lines selected for resistance to mitoxantrone or anthracyclines. ABCG2 consists of a nucleotide-binding domain (NBD) at the amino terminus and a transmembrane domain (TMD) at the carboxyl terminus and it is postulated to form a homodimer to perform its biological functions. Over-expression of ABCG2 in cell lines confers resistance on a wide variety of anticancer drugs including mitoxantrone,



daunorubicin, doxorubicin, **topotecan** and epirubicin. The expression of ABCG2 has been implicated in multidrug resistance (MDR) of acute myeloid leukemia and some solid tumors. In addition, ABCG2 can transport several fluorescent dyes or toxins. ABCG2 is found to be expressed in epithelial cells of intestine and colon, liver canaliculi, and renal tubules, where it serves to eliminate the plasma level of orally administered anticancer drugs as well as ingested toxins. ABCG2 is found to be highly expressed in placenta and the luminal surface of microvessel endothelium blood-brain barrier where it may play a role in limiting the penetration of drugs, such as **topotecan** from the maternal plasma into the fetus and from blood to brain. A variety of inhibitors for ABCG2 including GF120918 may prove useful for sensitizing cancer cells to chemotherapy or altering the distribution of orally administered drug substrates of ABCG2. Interestingly, ABCG2 is also expressed highly in hematopoietic stem cells. However, the function of ABCG2 in stem cells is currently unknown, although it may provide protection to stem cells from a variety of xenobiotics.

CT

## Medical Descriptors:

- \*multidrug resistance
- \*cancer chemotherapy
- protein expression
- molecular cloning
- nucleotide binding site
- amino terminal sequence
- carboxy terminal sequence
- gene overexpression
- cell line
- acute granulocytic leukemia: DR, drug resistance
- solid tumor: DR, drug resistance
- intestine epithelium cell
- colon mucosa
- intrahepatic bile duct
- kidney tubule
- vascular endothelium
- blood brain barrier
- maternal plasma
- hematopoietic stem cell
- cell protection
- drug blood level
- drug penetration
- drug bioavailability**
- drug transport
- drug structure
- drug distribution
- drug mechanism
- drug potentiation
- human
- nonhuman
- review
- nucleotide sequence
- Drug Descriptors:
- \*xenobiotic agent: AN, drug analysis
- \*xenobiotic agent: CR, drug concentration
- \*xenobiotic agent: IT, drug interaction
- \*xenobiotic agent: PK, pharmacokinetics
- \*xenobiotic agent: PD, pharmacology
- \*xenobiotic agent: IV, intravenous drug administration
- \*xenobiotic agent: PO, oral drug administration
- \*breast cancer resistance protein: EC, endogenous compound
- placenta protein: EC, endogenous compound

mitoxantrone: PK, pharmacokinetics  
 anthracycline derivative: PK, pharmacokinetics  
 membrane protein: EC, endogenous compound  
 antineoplastic agent: AN, drug analysis  
 antineoplastic agent: CR, drug concentration  
 antineoplastic agent: PK, pharmacokinetics  
 antineoplastic agent: IV, intravenous drug administration  
 antineoplastic agent: PO, oral drug administration  
 daunorubicin: PK, pharmacokinetics  
 doxorubicin: PK, pharmacokinetics  
   **topotecan: IT, drug interaction**  
   **topotecan: PK, pharmacokinetics**  
   **topotecan: IV, intravenous drug administration**  
   **topotecan: PO, oral drug administration**  
   **elacridar: AN, drug analysis**  
   **elacridar: PK, pharmacokinetics**  
 epirubicin: PK, pharmacokinetics  
 fluorescent dye  
 toxin  
 prazosin: PK, pharmacokinetics  
 etoposide: PK, pharmacokinetics  
 teniposide: PK, pharmacokinetics  
   **irinotecan: PK, pharmacokinetics**  
   **9 aminocamptothecin: PK, pharmacokinetics**  
 7 ethyl 10 hydroxycamptothecin: IT, drug interaction  
 7 ethyl 10 hydroxycamptothecin: PK, pharmacokinetics  
 7 ethyl 10 hydroxycamptothecin: PD, pharmacology  
 6 formylamino 12,13 dihydro 1,11 dihydroxy 5h indolo[2,3 a]pyrrolo[3,4 c]carbazole 5,7(6h) dione 13 glucoside: PK, pharmacokinetics  
 cytotoxic agent: AN, drug analysis  
 cytotoxic agent: CR, drug concentration  
 cytotoxic agent: IT, drug interaction  
 cytotoxic agent: PK, pharmacokinetics  
 cytotoxic agent: PD, pharmacology  
 cytotoxic agent: IV, intravenous drug administration  
 cytotoxic agent: PO, oral drug administration  
 flavopiridol: PK, pharmacokinetics  
 bisantrene: PK, pharmacokinetics  
 methotrexate: PK, pharmacokinetics  
 zidovudine: PK, pharmacokinetics  
 pheophorbide: AN, drug analysis  
 pheophorbide: PK, pharmacokinetics  
 reserpine: AN, drug analysis  
 reserpine: PK, pharmacokinetics  
 reserpine: PD, pharmacology  
 tamoxifen: AN, drug analysis  
 tamoxifen: PK, pharmacokinetics  
 unindexed drug

RN (mitoxantrone) 65271-80-9, 70476-82-3; (daunorubicin) 12707-28-7, 20830-81-3, 23541-50-6; (doxorubicin) 23214-92-8, 25316-40-9; (**topotecan**) 119413-54-6, 123948-87-8; (elacridar) 143664-11-3; (epirubicin) 56390-09-1, 56420-45-2; (prazosin) 19216-56-9, 19237-84-4; (etoposide) 33419-42-0; (teniposide) 29767-20-2; (**irinotecan**) 100286-90-6; (7 ethyl 10 hydroxycamptothecin) 86639-52-3; (6 formylamino 12,13 dihydro 1,11 dihydroxy 5h indolo[2,3 a]pyrrolo[3,4 c]carbazole 5,7(6h) dione 13 glucoside) 151069-12-4; (flavopiridol) 146426-40-6; (bisantrene) 71439-68-4, 78186-34-2; (methotrexate) 15475-56-6, 59-05-2, 7413-34-5; (zidovudine) 30516-87-1; (reserpine) 50-55-5, 8001-95-4; (tamoxifen) 10540-29-1  
 CN **Gf 120918; Cpt 11; Sn 38; Nb 506**

GEN GENBANK AAA82056 referred number; GENBANK AAC97367 referred number;  
 GENBANK AAG40003 referred number; GENBANK AAG40004 referred number;  
 GENBANK AAK14241 referred number; GENBANK AAO13805 referred number;  
 GENBANK CAA36038 referred number; GENBANK CAA62631 referred number;  
 GENBANK NP\_036050 referred number; GENBANK NP\_071452 referred number;  
 GENBANK NP\_114090 referred number; GENBANK P12428 referred number; GENBANK  
 Q64343 referred number

L154 ANSWER 4 OF 25 WPIX COPYRIGHT 2004 THOMSON DERWENT on STN

AN 2003-381415 [36] WPIX

DNC C2003-101173

TI Mucoadhesive vaginal composition for treating cancer comprises  
 chemotherapeutic agent, lipophilic or hydrophilic carrier, mucoadhesive  
 agent and sorption promoter.

DC A96 B05 B07 P32

IN BENET, L Z; LIU, J H; PAULETTI, G M; RITSCHER, W A

PA (BENE-I) BENET L Z; (LIUJ-I) LIU J H; (PAUL-I) PAULETTI G M; (RITS-I)  
 RITSCHER W A; (UMDU-N) UMD INC

CYC 100

PI WO 2003020210 A2 20030313 (200336)\* EN 61 A61K000-00

RW: AT BE BG CH CY CZ DE DK EA EE ES FI FR GB GH GM GR IE IT KE LS LU

MC MW MZ NL OA PT SD SE SK SL SZ TR TZ UG ZM ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK

DM DZ EC EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR

KZ LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ OM PH PL PT

RO RU SD SE SG SI SK SL TJ TM TN TR TT TZ UA UG UZ VN YU ZA ZM ZW

US 2003049302 A1 20030313 (200336) A61K009-22

ADT WO 2003020210 A2 WO 2002-US27027 20020821; US 2003049302 A1 Provisional US  
 2001-315877P 20010829, US 2002-226667 20020821

PRAI US 2001-315877P 20010829; US 2002-226667 20020821

IC ICM A61K000-00; A61K009-22

ICS A61F006-06

AB WO2003020210 A UPAB: 20030609

NOVELTY - A mucoadhesive vaginal composition comprises a chemotherapeutic  
 agent, a lipophilic or hydrophilic carrier, a mucoadhesive agent and a  
 sorption promoter.

DETAILED DESCRIPTION - A mucoadhesive vaginal composition comprises:  
 a chemotherapeutic agent (0.001 - 3000 mg) (preferably daunorubicin,  
 doxorubicin, idarubicin, amrubicin, pirarubicin, epirubicin, mitoxantrone,  
 etoposide, teniposide, vinblastine, vincristine, mitomycin C, paclitaxel,  
 docetaxel, actinomycin D, colchicine, **topotecan**, irinotecan or  
 gemcitabine; or a membrane efflux system inhibitor (preferably  
 cyclosporin, verapamil, valspodar, biricodar, quinidine, terfenadine,  
 pervilleine A, **GF120918**, LY335979, OC144-093, **XR9576**,  
 probenecid and/or MK571); a lipophilic or hydrophilic carrier (30 - 95) %  
 (preferably saturated mono-, di- and/or triglyceride of 8-18C fatty acid,  
 polyethylene glycol (having molecular weight of 200 - 800)) and/or their  
 derivative; a mucoadhesive agent (5 - 25) % (preferably cellulose  
 derivative, natural gum, alginate or pectin); and a sorption promoter (5 -  
 25) % (preferably non-ionizable glycol ester derivative, glycol derivative  
 with glycerol ester, non-ionizable glycol ether derivative or an  
 inter-esterified stone oil).

An INDEPENDENT CLAIM is included for a medicated intravaginal device  
 for transmucosal delivery of chemotherapeutic agents and membrane efflux  
 system inhibitors comprising (C1).

ACTIVITY - Cytostatic.

MECHANISM OF ACTION - None given.

USE - For treating cancer in human females (claimed) (e.g. ovarian,  
 cervical and uterine cancer).

ADVANTAGE - The composition allows systemic circulation of the drug

and permits administration of lower concentrations of the drug. The composition prevents leaking of the drug out of the vagina, which results in greater systemic bioavailability compared to that after oral administration; and enhances transmucosal absorption of chemotherapeutic agents. The vaginal delivery of inhibitor of membrane efflux system further reduces the risk of toxic side effects following administration to cancer patients diagnosed with drug-resistant tumors. The method avoids intravenous administration, permits extended or controlled continuous or pulsed delivery of the chemotherapeutic agents and/or inhibitors of membrane efflux system and thus achieves delivery of higher concentration of the drug.

Dwg.1/17

FS CPI GMPI

FA AB; GI; DCN

MC CPI: A12-V01; B02-Z; B04-A02; B04-B03A; B04-C01; B04-C02A; B04-C02D; B04-C03C; B06-H; B07-H; B10-A08; B11-C04; B12-M02B; B12-M03; B12-M08; B12-M11C; B12-M11E; B12-M11F; B14-H01B

L154 ANSWER 5 OF 25 WPIX COPYRIGHT 2004 THOMSON DERWENT on STN

AN 2003-553619 [52] WPIX

DNC C2003-149550

TI New use of **acridine derivative** for inhibiting mitoxantrone resistance transporter in treatment of e.g. cancer, acute lymphocytic leukemia and non-Hodgkin's lymphoma.

DC B02 B04 D16

IN BATES, S; ROBEY, R

PA (USSH) US DEPT HEALTH & HUMAN SERVICES

CYC 1

PI US 6521635 B1 20030218 (200352)\* 27 A61K031-435

ADT US 6521635 B1 Provisional US 2000-177410P 20000120, US 2001-767594 20010122

PRAI US 2000-177410P 20000120; US 2001-767594 20010122

IC ICM A61K031-435

AB US 6521635 B UPAB: 20030813

NOVELTY - Use of an **acridine derivative** (I), or its salt or solvate, for inhibiting a mitoxantrone resistance (MXR) transporter in a cell overexpressing MXR gene but not overexpressing a P-glycoprotein (Pgp) gene, by contacting with the cell, is new.

DETAILED DESCRIPTION - Use of an **acridine derivative** of formula (I), or its salt or solvate, for inhibiting a mitoxantrone resistance (MXR) transporter in a cell overexpressing MXR gene but not overexpressing a P-glycoprotein (Pgp) gene, by contacting with the cell, is new.

Ra = R1, amino or nitro;

R1 = H, halo, 1-4C alkyl, 1-4C alkoxy or 1-4C alkylthio;

R2 = H or 1-4C alkyl;

A = O, S or a bond;

B' = unsubstituted 1-4C alkylene; and

R3, R4 = 1-4C alkoxy.

INDEPENDENT CLAIMS are also included for the following:

(1) assaying the modulation of the functional effect of a test compound on the cell by (I), comprising contacting the test compound with the cells in the presence and absence of (I) and measuring the ability of (I) to modulate the functional effect of the test compound; and

(2) treatment of cancer that overexpresses the (MXR) gene but not overexpresses the (Pgp) gene, comprising co-administering a chemotherapeutic recognized by the MXR transporter and (I).

ACTIVITY - Cytostatic.

MECHANISM OF ACTION - MXR transporter inhibitor; Pgp antagonist; Multidrug resistance gene 1 (MDR-1)-inhibitor; Cancer cell growth

inhibitor.

Colon cancer cell line (S1-M1-80) (2000 cells/well), mitoxantrone and N-(4-(2-1,2,3,4-tetrahydro-6,7-dimethoxy-2-isoquinoliny1)-ethyl)-phenyl)-9,10-dihydro-5-methoxy-9-oxo-4-acridine carboxamide (A) were added to a 96-well plate. Control was prepared without (A), and it was incubated at 37 deg. C for 96 hours. After incubation, the cells were fixed and stained with sulforhodamine B, and IC50 was determined. The IC50 (micro M) of mitoxantrone, without and with (A) (1 micro M), was 222 plus or minus 101 and 0.12 plus or minus 0.11, respectively. The results showed that (A) sensitized S1-M1-80 cells expressing MXR by 1850-fold.

USE - For inhibiting the MXR transporter, and for treating cancer (e.g. lung cancer, colon cancer, breast cancer, prostate cancer, acute lymphocytic leukemia, non-Hodgkin's lymphoma and ovarian cancer) (claimed).

ADVANTAGE - (I) Is a potent multispecific antagonist capable of inhibiting or reversing both Pgp-mediated and MXR-mediate multidrug resistance phenotype.

Dwg.0/2

FS

CPI

FA

AB; GI; DCN

MC

CPI: B02-D; B04-F01; B06-D03; B06-D11; B06-E05; B08-D02; B11-C08E; B12-K04E; B14-H01; B14-L06; D05-H09

L154 ANSWER 6 OF 25 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.  
on STN

AN

2003195141 EMBASE

TI

Pharmacogenomics of ABC transporters and its role in cancer chemotherapy.

AU

Sparreboom A.; Danesi R.; Ando Y.; Chan J.; Figg W.D.

CS

W.D. Figg, Clinical Pharmacology Research Core, Center for Cancer Research, National Cancer Institute, 9000 Rockville Pike, Bethesda, MD 20892, United States. wdfigg@helix.nih.gov

SO

Drug Resistance Updates, (2003) 6/2 (71-84).

Refs: 135

ISSN: 1368-7646 CODEN: DRUPFW

CY

United Kingdom

DT

Journal; (Short Survey)

FS

016 Cancer

022 Human Genetics

029 Clinical Biochemistry

030 Pharmacology

037 Drug Literature Index

LA

English

SL

English

AB

ATP-binding cassette (ABC) genes play a role in the resistance of malignant cells to anticancer agents. The ABC gene products, including ABCB1 (P-glycoprotein), ABCC1 (MRP1), ABCC2 (MRP2, cMOAT), and ABCG2 (BCRP, MXR, ABCP) are also known to influence oral absorption and disposition of a wide variety of drugs. As a result, the expression levels of these proteins in humans have important consequences for an individual's susceptibility to certain drug-induced side effects, interactions, and treatment efficacy. Naturally occurring variants in ABC transporter genes have been identified that might affect the function and expression of the protein. This review focuses on recent advances in the pharmacogenomics of ABC transporters, and discusses potential implications of genetic variants for the chemotherapeutic treatment of cancer.

CT

Medical Descriptors:

\*pharmacogenomics

\*cancer chemotherapy

\*multidrug resistance

cancer cell

drug absorption  
drug disposition  
protein expression  
protein blood level  
drug sensitivity  
drug efficacy  
protein function  
genetic variability  
protein motif  
protein domain  
genetic code  
gene mapping  
chromosome 7q  
phenotype  
drug transport  
blood brain barrier  
cell type  
epithelium cell  
**drug bioavailability**  
chromosome 16p  
drug conjugation  
knockout mouse  
drug hypersensitivity  
dendritic cell  
chromosome 10q  
gene mutation  
anion transport  
protein localization  
gene identification  
gene function  
detoxification  
genetic screening  
gene sequence  
wild type  
exon  
missense mutation  
genetic analysis  
Dubin Johnson syndrome  
genotype  
drug metabolism  
oxidative stress  
rhabdomyosarcoma: DR, drug resistance  
human  
nonhuman  
short survey  
priority journal  
Drug Descriptors:  
\*antineoplastic agent: AD, drug administration  
\*antineoplastic agent: PO, oral drug administration  
\*gene product: EC, endogenous compound  
\*ABC transporter: EC, endogenous compound  
glycoprotein P: EC, endogenous compound  
multidrug resistance protein 1: EC, endogenous compound  
multidrug resistance protein 2: EC, endogenous compound  
glutathione  
etoposide: TO, drug toxicity  
leukotriene C4: EC, endogenous compound  
radixin: EC, endogenous compound  
mitoxantrone  
anthracycline

**topotecan**

doxorubicin

protein: EC, endogenous compound

breast cancer resistance protein: EC, endogenous compound

mitoxantrone resistance protein: EC, endogenous compound

dye

digoxin: CR, drug concentration

digoxin: IV, intravenous drug administration

digoxin: PO, oral drug administration

tsukubaenolide: CR, drug concentration

cyclosporin

vincristine

valspodar: PD, pharmacology

biricodar: PD, pharmacology

dactinomycin

daunorubicin

cyclosporin A

chlorambucil

clotrimazole

unindexed drug

unclassified drug

**xr 9576**

- RN (multidrug resistance protein 2) 256503-65-8; (glutathione) 70-18-8;  
 (etoposide) 33419-42-0; (leukotriene C4) 72025-60-6; (mitoxantrone)  
 65271-80-9, 70476-82-3; (**topotecan**) 119413-54-6, 123948-87-8;  
 (doxorubicin) 23214-92-8, 25316-40-9; (protein) 67254-75-5; (digoxin)  
 20830-75-5, 57285-89-9; (tsukubaenolide) 104987-11-3; (cyclosporin)  
 79217-60-0; (vincristine) 57-22-7; (valspodar) 121584-18-7; (biricodar)  
 174254-13-8; (dactinomycin) 1402-38-6, 1402-58-0, 50-76-0; (daunorubicin)  
 12707-28-7, 20830-81-3, 23541-50-6; (cyclosporin A) 59865-13-3,  
 63798-73-2; (chlorambucil) 305-03-3; (clotrimazole) 23593-75-1
- CN **Xr 9576**

L154 ANSWER 7 OF 25 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.  
 on STN

AN 2003030753 EMBASE

TI Mammalian drug efflux transporters of the ATP binding cassette (ABC)  
 family: An overview.

AU Schinkel A.H.; Jonker J.W.

CS A.H. Schinkel, Division of Experimental Therapy, Netherlands Cancer  
 Institute, Plesmanlaan 121, 1066 CX Amsterdam, Netherlands.  
 a.schinkel@nki.nl

SO Advanced Drug Delivery Reviews, (21 Jan 2003) 55/1 (3-29).

Refs: 169

ISSN: 0169-409X CODEN: ADDREP

PUI S 0169-409X(02)00169-2

CY Netherlands

DT Journal; General Review

FS 016 Cancer

030 Pharmacology

037 Drug Literature Index

038 Adverse Reactions Titles

LA English

SL English

AB Active drug efflux transporters of the ATP binding cassette  
 (ABC)-containing family of proteins have a major impact on the  
 pharmacological behavior of most of the drugs in use today.  
 Pharmacological properties affected by ABC transporters include the oral  
 bioavailability, hepatobiliary, direct intestinal, and urinary excretion  
 of drugs and drug-metabolites and -conjugates. Moreover, the penetration

of drugs into a range of important pharmacological sanctuaries, such as brain, testis, and fetus, and the penetration into specific cell- and tissue compartments can be extensively limited by ABC transporters. These interactions with ABC transporters determine to a large extent the clinical usefulness, side effects and toxicity risks of drugs. Many other xenotoxins, (pre-)carcinogens and endogenous compounds are also influenced by the ABC transporters, with corresponding consequences for the well-being of the individual. We aim to provide an overview of properties of the mammalian ABC transporters known to mediate significant transport of clinically relevant drugs. .COPYRGHT. 2002 Elsevier Science B.V. All rights reserved.

CT Medical Descriptors:  
 drug transport  
 protein family  
     **drug bioavailability**  
 hepatobiliary system  
 intestine absorption  
 urinary excretion  
 drug excretion  
 drug penetration  
 drug conjugation  
 blood brain barrier  
 fetomaternal transfusion  
 toxicity  
 risk factor  
 protein function  
 blood testis barrier  
 kidney  
 tissue distribution  
 protein localization  
 drug degradation  
 ataxia: SI, side effect  
 transport kinetics  
 human  
 nonhuman  
 clinical trial  
 review

priority journal  
 Drug Descriptors:  
 \*ABC transporter: EC, endogenous compound  
 breast cancer resistance protein: EC, endogenous compound  
 multidrug resistance protein 1: EC, endogenous compound  
 multidrug resistance protein 2: EC, endogenous compound  
 multidrug resistance protein 3: EC, endogenous compound  
 multidrug resistance protein 4: EC, endogenous compound  
 multidrug resistance protein 5: EC, endogenous compound  
 glycoprotein P: EC, endogenous compound  
 asimadoline  
 morphine  
 vinblastine  
 vincristine  
 paclitaxel  
 docetaxel: PK, pharmacokinetics  
 docetaxel: IV, intravenous drug administration  
 doxorubicin: DO, drug dose  
 doxorubicin: IT, drug interaction  
 doxorubicin: PK, pharmacokinetics  
 doxorubicin: IP, intraperitoneal drug administration  
 daunorubicin  
 epirubicin



bisantrene  
 mitoxantrone  
 etoposide: DO, drug dose  
 etoposide: IT, drug interaction  
 etoposide: PK, pharmacokinetics  
 etoposide: IP, intraperitoneal drug administration  
 valspodar: AE, adverse drug reaction  
 valspodar: CT, clinical trial  
 valspodar: DO, drug dose  
 valspodar: IT, drug interaction  
 valspodar: PK, pharmacokinetics  
 valspodar: PO, oral drug administration  
   elacridar: CT, clinical trial  
   elacridar: CM, drug comparison  
   elacridar: DO, drug dose  
   elacridar: IT, drug interaction  
   elacridar: PK, pharmacokinetics  
   elacridar: PD, pharmacology  
   elacridar: PO, oral drug administration  
 1 [4 (11,11 difluorodibenzo[b,e]bicyclo[5.1.0]oct 5 yl) 1 piperazinyl] 3  
 (5 quinolinyloxy) 2 propanol: CT, clinical trial  
 1 [4 (11,11 difluorodibenzo[b,e]bicyclo[5.1.0]oct 5 yl) 1 piperazinyl] 3  
 (5 quinolinyloxy) 2 propanol: CM, drug comparison  
 1 [4 (11,11 difluorodibenzo[b,e]bicyclo[5.1.0]oct 5 yl) 1 piperazinyl] 3  
 (5 quinolinyloxy) 2 propanol: IT, drug interaction  
 1 [4 (11,11 difluorodibenzo[b,e]bicyclo[5.1.0]oct 5 yl) 1 piperazinyl] 3  
 (5 quinolinyloxy) 2 propanol: PK, pharmacokinetics  
 1 [4 (11,11 difluorodibenzo[b,e]bicyclo[5.1.0]oct 5 yl) 1 piperazinyl] 3  
 (5 quinolinyloxy) 2 propanol: IP, intraperitoneal drug administration  
 1 [4 (11,11 difluorodibenzo[b,e]bicyclo[5.1.0]oct 5 yl) 1 piperazinyl] 3  
 (5 quinolinyloxy) 2 propanol: IV, intravenous drug administration  
 1 [4 (11,11 difluorodibenzo[b,e]bicyclo[5.1.0]oct 5 yl) 1 piperazinyl] 3  
 (5 quinolinyloxy) 2 propanol: PO, oral drug administration  
 protein inhibitor: AE, adverse drug reaction  
 protein inhibitor: CT, clinical trial  
 protein inhibitor: DO, drug dose  
 protein inhibitor: IT, drug interaction  
 protein inhibitor: PK, pharmacokinetics  
 protein inhibitor: IV, intravenous drug administration  
 protein inhibitor: PO, oral drug administration  
   xr 9576: IT, drug interaction  
   xr 9576: IV, intravenous drug administration  
   xr 9576: PO, oral drug administration  
 oc 144 093: IT, drug interaction  
 oc 144 093: IV, intravenous drug administration  
 oc 144 093: PO, oral drug administration  
 saquinavir: IT, drug interaction  
   topotecan  
 7 ethyl 10 hydroxycamptothecin  
 unindexed drug  
 unclassified drug  
 RN (multidrug resistance protein 2) 256503-65-8; (multidrug resistance  
 protein 3) 231947-64-1; (multidrug resistance protein 4) 299244-49-8;  
 (multidrug resistance protein 5) 266988-95-8; (asimadoline) 153205-46-0;  
 (morphine) 52-26-6, 57-27-2; (vinblastine) 865-21-4; (vincristine)  
 57-22-7; (paclitaxel) 33069-62-4; (docetaxel) 114977-28-5; (doxorubicin)  
 23214-92-8, 25316-40-9; (daunorubicin) 12707-28-7, 20830-81-3, 23541-50-6;  
 (epirubicin) 56390-09-1, 56420-45-2; (bisantrene) 71439-68-4, 78186-34-2;  
 (mitoxantrone) 65271-80-9, 70476-82-3; (etoposide) 33419-42-0; (valspodar)  
 121584-18-7; (elacridar) 143664-11-3; (1 [4 (11,11

difluorodibenzo[b,e]bicyclo[5.1.0]oct 5 yl) 1 piperazinyl] 3 (5 quinolinyl) 2 propanol) 167465-36-3; (saquinavir) 127779-20-8, 149845-06-7; (**topotecan**) 119413-54-6, 123948-87-8; (7 ethyl 10 hydroxycamptothecin) 86639-52-3

CN Sdz psc 833; **Gf 120918**; Ly 335979; Sn 38; Oc 144 093

L154 ANSWER 8 OF 25 MEDLINE on STN

AN 2002346178 MEDLINE

DN PubMed ID: 12089219

TI Boosting bioavailability of topotecan: what do we gain?.

CM Comment on: J Clin Oncol. 2002 Jul 1;20(13):2943-50. PubMed ID: 12089223

Comment in: J Clin Oncol. 2003 Jan 1;21(1):177; author reply 177. PubMed ID: 12506192

AU Hudes Gary

SO Journal of clinical oncology : official journal of the American Society of Clinical Oncology, (2002 Jul 1) 20 (13) 2918-9.

Journal code: 8309333. ISSN: 0732-183X.

CY United States

DT Commentary

Editorial

LA English

FS Priority Journals

EM 200207

ED Entered STN: 20020629

Last Updated on STN: 20030116

Entered Medline: 20020715

CT Check Tags: Female; Human

\*ATP-Binding Cassette Transporters: AI, antagonists & inhibitors

\*Acridines: PD, pharmacology

Administration, Oral

Antineoplastic Agents: AD, administration & dosage

\*Antineoplastic Agents: PK, pharmacokinetics

Biological Availability

Breast Neoplasms: DT, drug therapy

\*Breast Neoplasms: ME, metabolism

\*DNA Topoisomerases, Type I: AI, antagonists & inhibitors

Drug Resistance, Multiple

Drug Resistance, Neoplasm

Enzyme Inhibitors: AD, administration & dosage

\*Enzyme Inhibitors: PK, pharmacokinetics

Intestinal Absorption: DE, drug effects

\*Isoquinolines: PD, pharmacology

Neoplasm Proteins: AI, antagonists & inhibitors

\*P-Glycoprotein: AI, antagonists & inhibitors

\*Tetrahydroisoquinolines

Topotecan: AD, administration & dosage

\*Topotecan: PK, pharmacokinetics

RN **123948-87-8 (Topotecan); 143664-11-3 (GF 120918)**

CN 0 (ABCG2 protein, human); 0 (ATP-Binding Cassette Transporters); 0

(Acridines); 0 (Antineoplastic Agents); 0 (Enzyme Inhibitors); 0

(Isoquinolines); 0 (Neoplasm Proteins); 0 (P-Glycoprotein); 0

(Tetrahydroisoquinolines); EC 5.99.1.2 (DNA Topoisomerases, Type I)

L154 ANSWER 9 OF 25 ZCAPLUS COPYRIGHT 2004 ACS on STN

AN 2002:696659 ZCAPLUS

DN 137:222100

ED Entered STN: 13 Sep 2002


TI Improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and oral drug delivery compositions

IN Schellens, Johannes Henricus Matthias; Schinkel, Alfred Hermanus

PA Netherlands Cancer Institute, Neth.  
 SO U.S. Pat. Appl. Publ., 11 pp., Cont.-in-part of Appl. No. PCT/NL00/00331.  
 CODEN: USXXCO  
 DT Patent  
 LA English  
 IC ICM A61K031-517  
 ICS A61K031-4745; A61K031-473; A61K031-47; A61K031-12  
 NCL 514297000  
 CC 63-6 (Pharmaceuticals)  
 Section cross-reference(s): 1  
 FAN.CNT 2

|      | PATENT NO.  | KIND | DATE     | APPLICATION NO. | DATE         |
|------|---|------|----------|-----------------|--------------|
| PI   | US 2002128282   | A1   | 20020912 | US 2001-988285  | 20011119 <-- |
|      | WO 2000069390   | A2   | 20001123 | WO 2000-NL331   | 20000517 <-- |
|      | WO 2000069390   | A3   | 20011213 |                 |              |
|      | W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM |      |          |                 |              |
|      | RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG  |      |          |                 |              |
| PRAI | NL 1999-1012066   | A    | 19990517 | <--             |              |
|      | NL 1999-1012481   | A    | 19990630 | <--             |              |
|      | WO 2000-NL331   | A2   | 20000517 |                 |              |

GI



I

AB A method for increasing the systemic exposure of cells selected from tumor cells and normal cells to an orally administered pharmaceutically active compound, wherein a bioenhancer comprising an inhibitor of BCRP (breast cancer resistance protein) is orally administered concomitantly with said orally administered pharmaceutically active compound, and in which method the inhibitor is administered simultaneously with the pharmaceutical compound. Coadministration of a single oral dose of GF120918 (I) results in a profoundly increased systemic exposure to oral topotecan.

ST drug delivery oral bioavailability enhancer

IT Multidrug resistance proteins  
 RL: BSU (Biological study, unclassified); BIOL (Biological study) (BCRP (breast cancer resistance protein); improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and oral drug delivery compns.)

IT Antitumor agents  
 Drug bioavailability  
 Human

- (improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and oral drug delivery comps.)
- IT Mycotoxins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and oral drug delivery comps.)
- IT Drug delivery systems  
(oral; improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and oral drug delivery comps.)
- IT 56-65-5, Atp, biological studies  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(binding to; improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and oral drug delivery comps.)
- IT 123948-87-8, Topotecan 143664-11-3, GF120918  
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and oral drug delivery comps.)
- IT 84-65-1D, Anthraquinone, derivs. 19216-56-9, Prazosin 65271-80-9, Mitoxantrone 86639-52-3, Sn38 91421-42-0, 9-Nitrocamptothecin 91421-43-1, 9-Aminocamptothecin 100286-90-6, Cpt11 118974-02-0, Fumitremorgin C 149882-10-0, GG211 169869-90-3, DX8951f 180422-22-4, XR 9051 203923-89-1, BNP1350 206873-63-4, XR 9576  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and oral drug delivery comps.)

L154 ANSWER 10 OF 25 MEDLINE on STN DUPLICATE 1  
AN 2002346182 MEDLINE  
DN PubMed ID: 12089223  
TI Increased oral bioavailability of topotecan in combination with the breast cancer resistance protein and P-glycoprotein inhibitor GF120918.  
CM Comment in: J Clin Oncol. 2002 Jul 1;20(13):2918-9. PubMed ID: 12089219  
AU Kruijtzter C M F; Beijnen J H; Rosing H; ten Bokkel Huinink W W; Schot M; Jewell R C; Paul E M; Schellens J H M  
CS Department of Medical Oncology, the Netherlands Cancer Institute/Antoni van Leeuwenhoek Hospital, Amsterdam, the Netherlands.  
SO Journal of clinical oncology : official journal of the American Society of Clinical Oncology, (2002 Jul 1) 20 (13) 2943-50.  
Journal code: 8309333. ISSN: 0732-183X.  
CY United States  
DT (CLINICAL TRIAL)  
Journal; Article; (JOURNAL ARTICLE)  
(RANDOMIZED CONTROLLED TRIAL)  
LA English  
FS Priority Journals  
EM 200207  
ED Entered STN: 20020629  
Last Updated on STN: 20020716  
Entered Medline: 20020715  
AB PURPOSE: We discovered that breast cancer resistance protein (BCRP), a recently identified adenosine triphosphate-binding cassette drug transporter, substantially limits the oral bioavailability of topotecan in mdrla/1b(-/-) P-glycoprotein (P-gp) knockout and wild-type mice. GF120918 is a potent inhibitor of BCRP and P-gp. The aim was to increase the bioavailability of topotecan by GF120918. PATIENTS AND METHODS: In cohort A, eight patients received 1.0 mg/m(2) oral topotecan with or without

coadministration of one single oral dose of 1,000 mg GF120918 (day 1 or day 8). In cohort B, eight other patients received 1.0 mg/m<sup>2</sup> intravenous topotecan with or without 1,000 mg oral GF120918 to study the effect of GF120918 on the systemic clearance of topotecan. RESULTS: After oral topotecan, the mean area under the plasma concentration-time curve (AUC) of total topotecan increased significantly from 32.4 +/- 9.6 microg.h/L without GF120918 to 78.7 +/- 20.6 microg.h/L when GF120918 was coadministered (P = .008). The mean maximum plasma concentration of total topotecan increased from 4.1 +/- 1.5 microg/L without GF120918 to 11.5 +/- 2.4 microg/L with GF120918 (P = .008). The apparent bioavailability in this cohort increased significantly from 40.0% (range, 32% to 47%) to 97.1% (range, 91% to 120%) (P = .008). Interpatient variability of the apparent bioavailability was 17% without and 11% with GF120918. After intravenous administration of topotecan, coadministration of oral GF120918 had a small but statistically significant effect on the AUC and systemic clearance of total topotecan but no statistically significant effect on maximum plasma concentration and terminal half-life of total topotecan. CONCLUSION: Coadministration of the BCRP and P-gp inhibitor GF120918 resulted in a significant increase of the systemic exposure of oral topotecan. The apparent oral bioavailability increased from 40.0% without to 97.1% with GF120918.

CT Check Tags: Female; Human  
 \*ATP-Binding Cassette Transporters: AI, antagonists & inhibitors  
 Acridines: PD, pharmacology  
 \*Acridines: TU, therapeutic use  
 Administration, Oral  
 Adult  
 Antineoplastic Agents: AD, administration & dosage  
 \*Antineoplastic Agents: PK, pharmacokinetics  
 Biological Availability  
 \*Breast Neoplasms: DT, drug therapy  
 \*Breast Neoplasms: ME, metabolism  
 Drug Administration Schedule  
 Drug Resistance, Multiple  
 Drug Resistance, Neoplasm  
 Enzyme Inhibitors: AD, administration & dosage  
 \*Enzyme Inhibitors: PK, pharmacokinetics  
 Isoquinolines: PD, pharmacology  
 \*Isoquinolines: TU, therapeutic use  
 Middle Aged  
 Neoplasm Proteins: AI, antagonists & inhibitors  
 \*P-Glycoprotein: AI, antagonists & inhibitors  
 \*Tetrahydroisoquinolines  
 Topotecan: AD, administration & dosage  
 \*Topotecan: PK, pharmacokinetics  
 RN 123948-87-8 (Topotecan); 143664-11-3 (GF 120918)  
 CN 0 (ABCG2 protein, human); 0 (ATP-Binding Cassette Transporters); 0 (Acridines); 0 (Antineoplastic Agents); 0 (Enzyme Inhibitors); 0 (Isoquinolines); 0 (Neoplasm Proteins); 0 (P-Glycoprotein); 0 (Tetrahydroisoquinolines)

L154 ANSWER 11 OF 25 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.  
 on STN  
 AN 2002410079 EMBASE  
 TI Oral cancer treatment: Developments in chemotherapy and beyond.  
 AU O'Neill V.J.; Twelves C.J.  
 CS C.J. Twelves, Cancer Res. UK Dept. of Med. Oncol., University of Glasgow, Alexander Stone Building, Switchback Road, Glasgow G6I IBD, United Kingdom. c.twelves@beatson.gla.ac.uk  
 SO British Journal of Cancer, (21 Oct 2002) 87/9 (933-937).

Refs: 24  
 ISSN: 0007-0920 CODEN: BJCAAI  
 CY United Kingdom  
 DT Journal; General Review  
 FS 016 Cancer  
 030 Pharmacology  
 037 Drug Literature Index  
 038 Adverse Reactions Titles  
 039 Pharmacy  
 LA English  
 SL English  
 AB Oncology is one of the few areas of medicine where most patients are treated intravenously rather than receiving oral drugs. Recently, several oral anti-cancer drugs have been approved and there are many more in development. Oral chemotherapy is attractive because of its convenience and ease of administration, particularly in the palliative setting. With an increasing number of oral agents emerging, we can expect to see a rapid rise in the use of oral chemotherapy in years to come. This article reviews recent developments in oral chemotherapy, both of traditional cytotoxics and novel, targeted agents, from the viewpoint of patients, physicians, drug developers and health-care providers. .COPYRGT. 2002 Cancer Research UK.  
 CT Medical Descriptors:  
 \*cancer combination chemotherapy  
 \*mouth cancer: DM, disease management  
 \*mouth cancer: DT, drug therapy  
 drug approval  
 cancer palliative therapy  
 drug formulation  
 lung small cell cancer: DT, drug therapy  
 drug efficacy  
 diarrhea: SI, side effect  
 bone marrow suppression: SI, side effect  
 toxicity: SI, side effect  
 drug hypersensitivity: SI, side effect  
 antineoplastic activity  
 drug activity  
 drug absorption  
 cardiotoxicity: SI, side effect  
**drug bioavailability**  
 rash: SI, side effect  
 acne: SI, side effect  
 nausea: SI, side effect  
 drug targeting  
 molecular mechanics  
 drug infusion  
 vomiting: SI, side effect  
 patient compliance  
 health care system  
 health care cost  
 drug cost  
 human  
 clinical trial  
 review  
 priority journal  
 Drug Descriptors:  
 \*antineoplastic agent: AE, adverse drug reaction  
 \*antineoplastic agent: CT, clinical trial  
 \*antineoplastic agent: AD, drug administration  
 \*antineoplastic agent: CB, drug combination

\*antineoplastic agent: IT, drug interaction  
\*antineoplastic agent: DT, drug therapy  
\*antineoplastic agent: PE, pharmacoeconomics  
\*antineoplastic agent: PR, pharmaceuticals  
\*antineoplastic agent: PK, pharmacokinetics  
\*antineoplastic agent: PD, pharmacology  
\*antineoplastic agent: IV, intravenous drug administration  
\*antineoplastic agent: PO, oral drug administration  
cytotoxic agent: AE, adverse drug reaction  
cytotoxic agent: CT, clinical trial  
cytotoxic agent: AD, drug administration  
cytotoxic agent: CB, drug combination  
cytotoxic agent: IT, drug interaction  
cytotoxic agent: DT, drug therapy  
cytotoxic agent: PR, pharmaceuticals  
cytotoxic agent: PK, pharmacokinetics  
cytotoxic agent: PD, pharmacology  
cytotoxic agent: IV, intravenous drug administration  
cytotoxic agent: PO, oral drug administration  
topotecan: AE, adverse drug reaction  
topotecan: AD, drug administration  
topotecan: DT, drug therapy  
topotecan: IV, intravenous drug administration  
topotecan: PO, oral drug administration  
taxane derivative: AE, adverse drug reaction  
taxane derivative: CT, clinical trial  
taxane derivative: AD, drug administration  
taxane derivative: CB, drug combination  
taxane derivative: IT, drug interaction  
taxane derivative: DT, drug therapy  
taxane derivative: PK, pharmacokinetics  
taxane derivative: PD, pharmacology  
taxane derivative: IV, intravenous drug administration  
taxane derivative: PO, oral drug administration  
cyclosporin A: AE, adverse drug reaction  
cyclosporin A: CB, drug combination  
cyclosporin A: IT, drug interaction  
cyclosporin A: DT, drug therapy  
cyclosporin A: PD, pharmacology  
fluorouracil: AE, adverse drug reaction  
fluorouracil: AD, drug administration  
fluorouracil: CB, drug combination  
fluorouracil: DT, drug therapy  
fluorouracil: IV, intravenous drug administration  
fluorouracil: PO, oral drug administration  
capecitabine: DT, drug therapy  
capecitabine: PD, pharmacology  
tegafur: DT, drug therapy  
tegafur: PD, pharmacology  
5 ethynyluracil: AE, adverse drug reaction  
5 ethynyluracil: CT, clinical trial  
5 ethynyluracil: CB, drug combination  
5 ethynyluracil: IT, drug interaction  
5 ethynyluracil: DT, drug therapy  
5 ethynyluracil: PK, pharmacokinetics  
5 ethynyluracil: PD, pharmacology  
UFT: AE, adverse drug reaction  
UFT: CB, drug combination  
UFT: DT, drug therapy  
folinic acid: AE, adverse drug reaction

folinic acid: CB, drug combination  
folinic acid: DT, drug therapy  
fluoropyrimidine: CB, drug combination  
fluoropyrimidine: DV, drug development  
fluoropyrimidine: DT, drug therapy  
fluoropyrimidine: PD, pharmacology  
imatinib: DT, drug therapy  
imatinib: PK, pharmacokinetics  
imatinib: PD, pharmacology  
imatinib: PO, oral drug administration  
gefitinib: AE, adverse drug reaction  
gefitinib: DT, drug therapy  
gefitinib: PD, pharmacology  
gefitinib: PO, oral drug administration  
erlotinib: AE, adverse drug reaction  
erlotinib: DT, drug therapy  
erlotinib: PD, pharmacology  
erlotinib: PO, oral drug administration  
cetuximab: AE, adverse drug reaction  
cetuximab: DT, drug therapy  
cetuximab: IV, intravenous drug administration  
cyclophosphamide: AE, adverse drug reaction  
cyclophosphamide: DT, drug therapy  
cyclophosphamide: PO, oral drug administration  
bms 275183: DV, drug development  
bms 275183: DT, drug therapy  
bms 275183: PD, pharmacology  
bms 275183: PO, oral drug administration  
    camptothecin derivative: DV, drug development  
    camptothecin derivative: DT, drug therapy  
    camptothecin derivative: PK, pharmacokinetics  
    camptothecin derivative: PD, pharmacology  
    camptothecin derivative: PO, oral drug administration  
    elacridar: DV, drug development  
    elacridar: DT, drug therapy  
    elacridar: PD, pharmacology  
    elacridar: PO, oral drug administration  
2' cyano 2' deoxy 4 n palmitoylcytarabine: DV, drug development  
2' cyano 2' deoxy 4 n palmitoylcytarabine: DT, drug therapy  
2' cyano 2' deoxy 4 n palmitoylcytarabine: PD, pharmacology  
2' cyano 2' deoxy 4 n palmitoylcytarabine: PO, oral drug administration  
antimetabolite: DV, drug development  
antimetabolite: DT, drug therapy  
antimetabolite: PD, pharmacology  
antimetabolite: PO, oral drug administration  
protein tyrosine kinase inhibitor: DV, drug development  
protein tyrosine kinase inhibitor: DT, drug therapy  
protein tyrosine kinase inhibitor: PD, pharmacology  
protein tyrosine kinase inhibitor: PO, oral drug administration  
su 006668: DV, drug development  
su 006668: DT, drug therapy  
su 006668: PD, pharmacology  
su 006668: PO, oral drug administration  
cep 701: DV, drug development  
cep 701: DT, drug therapy  
cep 701: PD, pharmacology  
cep 701: PO, oral drug administration  
3 [(3,5 dimethyl 1h pyrrol 2 yl)methylene] 1,3 dihydro 2h indol 2 one: DV,  
drug development  
3 [(3,5 dimethyl 1h pyrrol 2 yl)methylene] 1,3 dihydro 2h indol 2 one: DT,



drug therapy  
 3 [(3,5 dimethyl 1h pyrrol 2 yl)methylene] 1,3 dihydro 2h indol 2 one: PD,  
 pharmacology  
 3 [(3,5 dimethyl 1h pyrrol 2 yl)methylene] 1,3 dihydro 2h indol 2 one: PO,  
 oral drug administration  
 tas 102: DV, drug development  
 tas 102: DT, drug therapy  
 tas 102: PD, pharmacology  
 tas 102: PO, oral drug administration  
 cp 461: DV, drug development  
 cp 461: DT, drug therapy  
 cp 461: PD, pharmacology  
 cp 461: PO, oral drug administration  
 pkc 412: DV, drug development  
 pkc 412: DT, drug therapy  
 pkc 412: PD, pharmacology  
 pkc 412: PO, oral drug administration  
 unindexed drug  
 unclassified drug  
 diflomotecan  
 bms 294662

RN (topotecan) 119413-54-6, 123948-87-8; (cyclosporin A)  
 59865-13-3, 63798-73-2; (fluorouracil) 51-21-8; (capecitabine)  
 154361-50-9; (tegafur) 17902-23-7; (5 ethynyluracil) 59989-18-3; (UFT)  
 74578-38-4; (folinic acid) 58-05-9, 68538-85-2; (fluoropyrimidine)  
 675-21-8; (imatinib) 152459-95-5, 220127-57-1; (gefitinib) 184475-35-2,  
 184475-55-6, 184475-56-7; (erlotinib) 183319-69-9; (cetuximab)  
 205923-56-4; (cyclophosphamide) 50-18-0; (elacridar) 143664-11-3; (2'  
 cyano 2' deoxy 4 n palmitoylcytarabine) 151823-14-2; (cep 701)  
 111358-88-4, 156256-78-9; (3 [(3,5 dimethyl 1h pyrrol 2 yl)methylene] 1,3  
 dihydro 2h indol 2 one) 186610-95-7; (diflomotecan) 220997-97-7  
 CN Bn 80915; **Gf 120918**; Bms 275183; Cs 682; Zd 1839; Osi 774; Su  
 006668; Bms 294662; Su 5416; Tas 102; Cp 461; Pkc 412

L154 ANSWER 12 OF 25 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.  
 on STN

AN 2002218265 EMBASE

TI [Natural protection for the body. Drug transporters in the  
 gastro-intestinal tract].

GENEESMIDDELTRANSPORTERS IN HET MAAGDARMKANAAL: NATUURLIJKE BESCHERMERS  
 VAN HET LICHAAM.

AU Jorritsma A.; Schinkel A.H.; Schellens J.H.M.; Beijnen J.H.

CS Dr. J.H. Beijnen, Apotheek Slotervaartziekenhuis, Antoni van  
 Leeuwenhoekhuis, Louwesweg 6, 1066 EC Amsterdam, Netherlands

SO Pharmaceutisch Weekblad, (21 Jun 2002) 137/25 (904-910).

Refs: 25

ISSN: 0031-6911 CODEN: PHWEAW

CY Netherlands

DT Journal; (Short Survey)

FS 030 Pharmacology

037 Drug Literature Index

LA Dutch

SL English

AB Multidrug resistance research has revealed an important role for drug  
 transporters. These transmembrane transporters also appeared to have a  
 function under normal physiological circumstances, where they protect the  
 body against potentially harmful substances. In the gut, some of the  
 transporters are able to prevent the uptake of drugs, which causes a low  
 oral bioavailability. P-glycoprotein is the most studied transporter and  
 many P-glycoprotein substrates have low oral bioavailability. However, the

bioavailability of substrates strongly increases in P-glycoprotein-deficient mice and also when a P-glycoprotein inhibitor is co-administered with the substrate in wild type animals. Another group of drug transporters are the 'multidrug resistance-associated proteins' (MRPs). MRP(1) and MRP(2) are present under normal physiological circumstances where they have a protective function. Their influence on the oral bioavailability of substrates has not been determined yet. The 'breast cancer resistance protein' (BCRP) is a recently identified drug transporter. Like P-glycoprotein, BCRP is able to influence the intestinal uptake of drugs. There is a strong overlap between substrates of BCRP and P-glycoprotein and it is therefore difficult to determine the exact influence of BCRP on oral bioavailability. In the future, this will be possible by using recently developed BCRP-deficient mice. The various drug transporters can be important factors in the intestinal uptake of substrates. It is important to expand our knowledge about these transporters and safe inhibitors in order to improve the oral bioavailability of affected drugs.

CT Medical Descriptors:

\*intestine absorption

\*drug transport

protection

multidrug resistance

drug uptake

**drug bioavailability**

protein localization

protein binding

nonhuman

short survey

Drug Descriptors:

glycoprotein P: EC, endogenous compound

multidrug resistance protein: EC, endogenous compound

cell protein: EC, endogenous compound

breast cancer resistance protein: EC, endogenous compound

antineoplastic agent: CB, drug combination

antineoplastic agent: PK, pharmacokinetics

antineoplastic agent: PO, oral drug administration

**topotecan: CB, drug combination**

**topotecan: PK, pharmacokinetics**

**topotecan: PO, oral drug administration**

**elacridar: CB, drug combination**

**elacridar: PK, pharmacokinetics**

**elacridar: PO, oral drug administration**

unclassified drug

RN (multidrug resistance protein) 149200-37-3, 208997-77-7; (**topotecan**) 119413-54-6, 123948-87-8; (elacridar) 143664-11-3

L154 ANSWER 13 OF 25 MEDLINE on STN

AN 2002475189 MEDLINE

DN PubMed ID: 12237778

TI Induction of breast cancer resistance protein by the **camptothecin derivative** DX-8951f is associated with minor reduction of antitumour activity.

AU van Hattum A H; Hoogsteen I J; Schluper H M M; Maliepaard M; Scheffer G L; Scheper R J; Kohlhagen G; Pommier Y; Pinedo H M; Boven E

CS Department of Medical Oncology, Vrije Universiteit Medical Center, De Boelelaan 1117, 1081 HV Amsterdam, The Netherlands.

SO British journal of cancer, (2002 Sep 9) 87 (6) 665-72.

Journal code: 0370635. ISSN: 0007-0920.

CY Scotland: United Kingdom

DT Journal; Article; (JOURNAL ARTICLE)

LA English  
FS Priority Journals  
EM 200210  
ED Entered STN: 20020919  
Last Updated on STN: 20021026  
Entered Medline: 20021024

AB DX-8951f (exatecan mesylate), a new water-soluble **derivative of camptothecin**, is currently being evaluated in phase II clinical trials. Resistance may be acquired when treating cancer patients with DX-8951f. Therefore, we selected a subline of the human ovarian cancer cell line A2780 for resistance against DX-8951f to investigate possible mechanisms of resistance. This DX-8951f-resistant subline, designated 2780DX8 (resistance factor=9.3), displayed a typical cross-resistance pattern including compounds, such as topotecan (resistance factor =34), SN-38 (resistance factor =47), mitoxantrone (resistance factor =59) and doxorubicin (resistance factor =2.9), which have previously been associated with the expression of breast cancer resistance protein. 2780DX8 cells did not show changes in the topoisomerase I gene, in topoisomerase I protein levels or catalytic activity. Overexpression of breast cancer resistance protein could be detected, both at the mRNA and protein level, while staining for Pgp, MRP1, or LRP was negative. GF120918, an inhibitor of breast cancer resistance protein, was able to reverse the DX-8951f-induced resistance in 2780DX8 cells. In vivo experiments in well-established 2780DX8 human tumour xenografts demonstrated that the growth inhibition induced by CPT-11 was more affected by breast cancer resistance protein expression than that of DX-8951f. These data indicate for the first time that DX-8951f is able to induce breast cancer resistance protein as a mechanism of resistance. Breast cancer resistance protein, however, results in only minor reduction of antitumour activity of DX-8951f which is an advantage over topotecan and CPT-11/SN-38.

CT Check Tags: Comparative Study; Female; Human  
\*ATP-Binding Cassette Transporters: ME, metabolism  
Acridines: PD, pharmacology  
Animals  
Antigens, CD: ME, metabolism  
Antineoplastic Agents, Phytogenic: AD, administration & dosage  
\*Antineoplastic Agents, Phytogenic: TU, therapeutic use  
Breast Neoplasms: DT, drug therapy  
Breast Neoplasms: ME, metabolism  
Camptothecin: AD, administration & dosage  
Camptothecin: AA, analogs & derivatives  
\*Camptothecin: TU, therapeutic use  
Cell Division: DE, drug effects  
Cell Division: PH, physiology  
DNA Topoisomerases, Type I: AI, antagonists & inhibitors  
DNA Topoisomerases, Type I: GE, genetics  
DNA Topoisomerases, Type I: ME, metabolism  
Drug Resistance, Neoplasm  
Immunoenzyme Techniques  
Isoquinolines: PD, pharmacology  
\*Membrane Glycoproteins  
Mice  
Mice, Nude  
Mutation  
Neoplasm Proteins: ME, metabolism  
\*Neoplasms, Experimental: DT, drug therapy  
Neoplasms, Experimental: PA, pathology  
\*Ovarian Neoplasms: DT, drug therapy  
P-Glycoprotein: ME, metabolism

## \*Tetrahydroisoquinolines

Tetrazolium Salts: DU, diagnostic use

Thiazoles: DU, diagnostic use

Tumor Cells, Cultured: CY, cytology

Tumor Cells, Cultured: DE, drug effects

RN 143664-11-3 (GF 120918); 147785-22-6 (CD9 antigen); 298-93-1  
(thiazolyl blue); 7689-03-4 (Camptothecin)

CN 0 (ABCG2 protein, human); 0 (ATP-Binding Cassette Transporters); 0  
(Acridines); 0 (Antigens, CD); 0 (Antineoplastic Agents, Phytogenic);  
0 (DX 8951); 0 (Isoquinolines); 0 (Membrane Glycoproteins); 0  
(Neoplasm Proteins); 0 (P-Glycoprotein); 0 (Tetrahydroisoquinolines); 0  
(Tetrazolium Salts); 0 (Thiazoles); EC 5.99.1.2 (DNA Topoisomerases, Type  
I)

L154 ANSWER 14 OF 25 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.  
on STN

AN 2003003508 EMBASE

TI Improvement of oral drug treatment by temporary inhibition of drug  
transporters and/or cytochrome P450 in the gastrointestinal tract and  
liver: An overview.

AU Kruijtz C.M.F.; Beijnen J.H.; Schellens J.H.M.

CS Dr. J.H.M. Schellens, Netherlands Cancer Institute, Department of Medical  
Oncology, Plesmanlaan 121, 1066 CX Amsterdam, Netherlands. jhm@nki.nl

SO Oncologist, (2002) 7/6 (516-530).

Refs: 172

ISSN: 1083-7159 CODEN: OCOLF6

CY United States

DT Journal; Article

FS 016 Cancer

030 Pharmacology

037 Drug Literature Index

LA English

SL English

AB The oral bioavailability of many cytotoxic drugs is low and/or highly  
variable. This can be caused by high affinity for drug transporters and  
activity of metabolic enzymes in the gastrointestinal tract and liver. In  
this review, we will describe the main involved drug transporters and  
metabolic enzymes and discuss novel methods to improve oral treatment of  
affected substrate drugs. Results of preclinical and clinical phase I and  
II studies will be discussed in which affected substrate drugs, such as  
paclitaxel, docetaxel, and **topotecan**, are given orally in  
combination with an inhibitor of drug transport or drug metabolism. Future  
randomized studies will, hopefully, confirm that this strategy for oral  
treatment is at least as equally effective and safe as standard  
intravenous administration of these drugs.

CT Medical Descriptors:

\*drug inhibition

\*drug transport

gastrointestinal tract

liver

drug bioavailability

drug receptor binding

drug absorption

area under the curve

drug blood level

drug elimination

article

priority journal

Drug Descriptors:

\*cytochrome P450

\*cytotoxic agent: IT, drug interaction  
 \*cytotoxic agent: PK, pharmacokinetics  
 \*cytotoxic agent: PD, pharmacology  
 \*cytotoxic agent: IV, intravenous drug administration  
 \*cytotoxic agent: PO, oral drug administration  
 \*paclitaxel: PK, pharmacokinetics  
 \*paclitaxel: PD, pharmacology  
 \*paclitaxel: IV, intravenous drug administration  
 \*paclitaxel: PO, oral drug administration  
 \*docetaxel: IT, drug interaction  
 \*docetaxel: PK, pharmacokinetics  
 \*docetaxel: PD, pharmacology  
 \*docetaxel: PO, oral drug administration  
 \*topotecan: PK, pharmacokinetics  
 \*topotecan: PD, pharmacology  
 \*topotecan: IV, intravenous drug administration  
 \*topotecan: PO, oral drug administration  
 verapamil: PD, pharmacology  
 quinidine: PD, pharmacology  
 cyclosporin A: PD, pharmacology  
 elacridar: PD, pharmacology  
 1 [4 (11,11 difluorodibenzo[b,e]bicyclo[5.1.0]oct 5 yl) 1 piperazinyl] 3  
 (5 quinolinyloxy) 2 propanol: PD, pharmacology  
 valspodar: PD, pharmacology  
 biricodar: PD, pharmacology  
 multidrug resistance protein  
 ritonavir: IT, drug interaction  
 irinotecan: IT, drug interaction  
 irinotecan: PK, pharmacokinetics  
 irinotecan: PD, pharmacology  
 ketoconazole: IT, drug interaction  
 (cytochrome P450) 9035-51-2; (paclitaxel) 33069-62-4; (docetaxel)  
 114977-28-5; (**topotecan**) 119413-54-6, 123948-87-8; (verapamil)  
 152-11-4, 52-53-9; (quinidine) 56-54-2; (cyclosporin A) 59865-13-3,  
 63798-73-2; (elacridar) 143664-11-3; (1 [4 (11,11  
 difluorodibenzo[b,e]bicyclo[5.1.0]oct 5 yl) 1 piperazinyl] 3 (5  
 quinolinyloxy) 2 propanol) 167465-36-3; (valspodar) 121584-18-7;  
 (biricodar) 174254-13-8; (multidrug resistance protein) 149200-37-3,  
 208997-77-7; (ritonavir) 155213-67-5; (**irinotecan**) 100286-90-6;  
 (ketoconazole) 65277-42-1

L154 ANSWER 15 OF 25 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.  
on STN

AN 2003283112 EMBASE

TI Pharmacokinetics in cancer treatment: Clinical implications of  
interindividual variability and drug interactions.

AU Boven E.

CS Dr. E. Boven, Department of Medical Oncology, Vrije Universiteit Medical  
Centre, De Boelelaan 1117, Amsterdam, 1081 HV, Netherlands.  
e.boven@vumc.nl

SO American Journal of Cancer, (2002) 1/1 (33-53).

Refs: 205

ISSN: 1175-6357 CODEN: AJCMCB

CY New Zealand

DT Journal; General Review

FS 016 Cancer

030 Pharmacology

037 Drug Literature Index

038 Adverse Reactions Titles

LA English

SL English

AB The design of chemotherapy schedules for treatment of malignancies is based on the selection of optimal drug doses with tolerable adverse effects. Interindividual variation in absorption, distribution, metabolism and excretion may exist for a given dose, which depend on both physiological and pathological factors. These factors will be of importance for the outcome of treatment in terms of efficacy as well as toxicity. As chemotherapy usually consists of a combination of drugs, pharmacological interactions may be expected. This also holds for other drug classes, such as analgesics, antidepressants, antibiotics, that are frequently coadministered to patients receiving chemotherapy. The study of pharmacokinetics can give insight in to the extent of interindividual variability based on genetic and patient factors, as well as in the occurrence of drug interactions. Many anticancer agents need specific enzymes for their metabolism. Polymorphisms in gene expression resulting in differences in enzyme activity have been described, such as thiopurine methyltransferase for metabolism of 6-mercaptopurine, dihydropyrimidine dehydrogenase for fluorouracil and uridine diphosphate (UDP) glucuronosyl transferase 1A1 for SN-38 (the active metabolite of **irinotecan**). Cytochrome P450 isoenzymes form a very important drug-metabolizing family and CYP3A4 is responsible for the metabolism of several classes of drugs. This isoenzyme system can easily be induced or inhibited by other drugs. Interactions combining drugs requiring both CYP3A4 for metabolism and P170-glycoprotein (Pgp) for transport may result in enhanced adverse effects in patients. A well-known example is the interference of taxanes with the pharmacokinetics of anthracyclines. Patient factors, other than variable expression of drug-metabolizing enzymes, that may account for altered pharmacokinetic properties are: age, obesity, hypoalbuminemia, impaired renal or liver function. A combination of these factors may occur, especially in patients with advanced cancer. The presence of a drug interaction may be advantageous in some instances. For example, the limited oral bioavailability of paclitaxel may be improved by inhibition of Pgp-mediated drug efflux from the intestine. The same holds true for blocking the breast cancer resistance protein transporter in the intestine to enhance oral absorption of **topotecan**. It is only through prospective, preclinical and early clinical evaluation of both pharmacokinetics and pharmacodynamics, i.e. the effects of the drug on the body, that the pharmacological behavior of a particular drug can be identified. Changes in drug dose, sequence, or infusion duration, increase of the time-interval between drugs, etc., can be measures required to provide an optimal therapeutic index of combination chemotherapy for the patient with cancer.

CT Medical Descriptors:

- \*cancer: DR, drug resistance
- \*cancer: DT, drug therapy
- \*advanced cancer: DT, drug therapy
- heredity
- cancer patient
- cancer chemotherapy
- drug tolerability
- drug choice
- drug absorption
- drug distribution
- drug metabolism
- drug excretion
- protein function
- gene expression
- DNA polymorphism
- enzyme activity
- drug transport

age  
obesity  
hypoalbuminemia  
kidney dysfunction  
liver dysfunction  
    **drug bioavailability**  
intestine absorption  
infusion  
neurotoxicity: DT, drug therapy  
neurotoxicity: SI, side effect  
gastrointestinal tumor: DT, drug therapy  
colorectal carcinoma: DT, drug therapy  
enzyme deficiency  
side effect: SI, side effect  
Crigler Najjar syndrome: SI, side effect  
Gilbert disease: SI, side effect  
diarrhea: SI, side effect  
neutropenia: SI, side effect  
mucosa inflammation: SI, side effect  
lymphatic leukemia: DT, drug therapy  
bone marrow suppression: SI, side effect  
multidrug resistance  
human  
clinical trial  
review  
priority journal  
Drug Descriptors:  
\*antineoplastic agent: AE, adverse drug reaction  
\*antineoplastic agent: CT, clinical trial  
\*antineoplastic agent: IT, drug interaction  
\*antineoplastic agent: DT, drug therapy  
\*antineoplastic agent: PK, pharmacokinetics  
\*anthracycline derivative: AE, adverse drug reaction  
\*anthracycline derivative: CT, clinical trial  
\*anthracycline derivative: IT, drug interaction  
\*anthracycline derivative: DT, drug therapy  
\*anthracycline derivative: PK, pharmacokinetics  
\*cyclophosphamide derivative: AE, adverse drug reaction  
\*cyclophosphamide derivative: CT, clinical trial  
\*cyclophosphamide derivative: IT, drug interaction  
\*cyclophosphamide derivative: DT, drug therapy  
\*cyclophosphamide derivative: PK, pharmacokinetics  
\*ifosfamide: CT, clinical trial  
\*ifosfamide: IT, drug interaction  
\*ifosfamide: DT, drug therapy  
\*ifosfamide: PK, pharmacokinetics  
\*epipodophyllotoxin derivative: CT, clinical trial  
\*epipodophyllotoxin derivative: IT, drug interaction  
\*epipodophyllotoxin derivative: DT, drug therapy  
\*epipodophyllotoxin derivative: PK, pharmacokinetics  
\*taxane derivative: CT, clinical trial  
\*taxane derivative: IT, drug interaction  
\*taxane derivative: DT, drug therapy  
\*taxane derivative: PK, pharmacokinetics  
Vinc alkaloid: CT, clinical trial  
Vinc alkaloid: IT, drug interaction  
Vinc alkaloid: DT, drug therapy  
Vinc alkaloid: PK, pharmacokinetics  
    **camptothecin derivative: CT, clinical trial**  
    **camptothecin derivative: IT, drug interaction**

camptothecin derivative: DT, drug therapy  
camptothecin derivative: PK, pharmacokinetics  
platinum derivative: CT, clinical trial  
platinum derivative: IT, drug interaction  
platinum derivative: DT, drug therapy  
platinum derivative: PK, pharmacokinetics  
gemcitabine: CT, clinical trial  
gemcitabine: IT, drug interaction  
gemcitabine: DT, drug therapy  
gemcitabine: PK, pharmacokinetics  
glycoprotein P  
glycoprotein p170  
verapamil: CT, clinical trial  
verapamil: IT, drug interaction  
verapamil: DT, drug therapy  
verapamil: PK, pharmacokinetics  
cyclosporin derivative: AE, adverse drug reaction  
cyclosporin derivative: CT, clinical trial  
cyclosporin derivative: IT, drug interaction  
cyclosporin derivative: DT, drug therapy  
cyclosporin derivative: PK, pharmacokinetics  
analgesic agent: IT, drug interaction  
antidepressant agent: IT, drug interaction  
antibiotic agent: IT, drug interaction  
thiopurine methyltransferase: EC, endogenous compound  
cytochrome P450 isoenzyme: EC, endogenous compound  
mercaptopurine: CT, clinical trial  
mercaptopurine: IT, drug interaction  
mercaptopurine: DT, drug therapy  
mercaptopurine: PK, pharmacokinetics  
dihydropyrimidine dehydrogenase: EC, endogenous compound  
fluorouracil: AE, adverse drug reaction  
fluorouracil: CT, clinical trial  
fluorouracil: IT, drug interaction  
fluorouracil: DT, drug therapy  
fluorouracil: PK, pharmacokinetics  
glucuronosyltransferase: EC, endogenous compound  
7 ethyl 10 hydroxycamptothecin: CT, clinical trial  
7 ethyl 10 hydroxycamptothecin: IT, drug interaction  
7 ethyl 10 hydroxycamptothecin: DT, drug therapy  
7 ethyl 10 hydroxycamptothecin: PK, pharmacokinetics  
irinotecan: AE, adverse drug reaction  
irinotecan: CT, clinical trial  
irinotecan: IT, drug interaction  
irinotecan: DT, drug therapy  
irinotecan: PK, pharmacokinetics  
cytochrome P450 3A4: EC, endogenous compound  
paclitaxel: CT, clinical trial  
paclitaxel: IT, drug interaction  
paclitaxel: DT, drug therapy  
paclitaxel: PK, pharmacokinetics  
paclitaxel: PO, oral drug administration  
carrier protein  
thymidine: DT, drug therapy  
unindexed drug  
unclassified drug  
2,4 bis(allylamino) 6 [4 [[2,2 bis(4 fluorophenyl)ethyl]amino]piperidino]  
1,3,5 triazine  
r 101933  
elacridar



RN (ifosfamide) 3778-73-2; (gemcitabine) 103882-84-4; (verapamil) 152-11-4, 52-53-9; (thiopurine methyltransferase) 67339-09-7; (mercaptapurine) 31441-78-8, 50-44-2, 6112-76-1; (dihydropyrimidine dehydrogenase) 9026-89-5; (fluorouracil) 51-21-8; (glucuronosyltransferase) 37329-64-9, 9030-08-4; (7 ethyl 10 hydroxycamptothecin) 86639-52-3; (**irinotecan**) 100286-90-6; (cytochrome P450 3A4) 329736-03-0; (paclitaxel) 33069-62-4; (carrier protein) 80700-39-6; (thymidine) 50-89-5; (2,4 bis(allylamino) 6 [4 [[2,2 bis(4 fluorophenyl)ethyl]aminolpiperidino] 1,3,5 triazine) 140945-01-3; (elacridar) 143664-11-3

CN Sn 38; S 9788; R 101933; **Gf 120918**

L154 ANSWER 16 OF 25 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.  
on STN

AN 2002013681 EMBASE

TI The use of oral cytotoxic and cytostatic drugs in cancer treatment.

AU Sparreboom A.; De Jonge M.J.A.; Verweij J.

CS A. Sparreboom, Department of Medical Oncology, Rotterdam Cancer Inst. (D.d.H.K.), University Hospital Rotterdam, 3075 EA Rotterdam, Netherlands. sparreboom@onch.azr.nl

SO European Journal of Cancer, (2002) 38/1 (18-22).

Refs: 28

ISSN: 0959-8049 CODEN: EJCAEL

PUI S 0959-8049(01)00322-7

CY United Kingdom

DT Journal; General Review

FS 010 Obstetrics and Gynecology

015 Chest Diseases, Thoracic Surgery and Tuberculosis

016 Cancer

037 Drug Literature Index

038 Adverse Reactions Titles

LA English

SL English

AB Although with a few exceptions, most new anticancer agents are initially developed for intravenous use, oral treatment with anticancer agents is, if feasible, to be preferred, as this route of administration is convenient to patients, reduces administration costs and facilitates the use of more chronic treatment regimens. Recent studies have identified various physiological barriers limiting the oral absorption of anticancer drugs. Presently, several strategies are explored to alter the low and variable oral bioavailability of several important anticancer agents by taking advantage of an intentional interaction between anticancer agents and drugs that modulate active intestinal drug transporters or (intestinal) enzymes. .COPYRGHT. 2002 Elsevier Science Ltd. All rights reserved.

CT Medical Descriptors:

\*ovary cancer: DT, drug therapy

\*lung small cell cancer: DT, drug therapy  
drug use

cancer therapy

cost effectiveness analysis

long term care

physiology

drug absorption

**drug bioavailability**

protein expression

area under the curve

dose response

drug potentiation

drug efficacy

drug safety  
 drug tolerability  
 patient compliance  
 outpatient department  
 blood toxicity: SI, side effect  
 human  
 nonhuman

Hypericum perforatum

review

priority journal

Drug Descriptors:

\*cytotoxic agent: AE, adverse drug reaction  
 \*cytotoxic agent: AD, drug administration  
 \*cytotoxic agent: CB, drug combination  
 \*cytotoxic agent: IT, drug interaction  
 \*cytotoxic agent: DT, drug therapy  
 \*cytotoxic agent: PK, pharmacokinetics  
 \*cytotoxic agent: IV, intravenous drug administration  
 \*cytotoxic agent: PO, oral drug administration  
 \*cytostatic agent: AE, adverse drug reaction  
 \*cytostatic agent: AD, drug administration  
 \*cytostatic agent: CB, drug combination  
 \*cytostatic agent: IT, drug interaction  
 \*cytostatic agent: DT, drug therapy  
 \*cytostatic agent: PK, pharmacokinetics  
 \*cytostatic agent: IV, intravenous drug administration  
 \*cytostatic agent: PO, oral drug administration

topotecan: AE, adverse drug reaction

topotecan: AD, drug administration

topotecan: DT, drug therapy

topotecan: PK, pharmacokinetics

topotecan: IV, intravenous drug administration

topotecan: PO, oral drug administration

intestine enzyme: EC, endogenous compound

glycoprotein P: EC, endogenous compound

protein: EC, endogenous compound

paclitaxel: AD, drug administration

paclitaxel: PK, pharmacokinetics

paclitaxel: PO, oral drug administration

taxane derivative: AD, drug administration

taxane derivative: CB, drug combination

taxane derivative: PK, pharmacokinetics

taxane derivative: PO, oral drug administration

cyclosporin A: AD, drug administration

cyclosporin A: CB, drug combination

cyclosporin A: IT, drug interaction

cyclosporin A: PK, pharmacokinetics

cyclosporin A: PO, oral drug administration

elacridar: AD, drug administration

elacridar: CB, drug combination

elacridar: PK, pharmacokinetics

elacridar: PO, oral drug administration

rifampicin: IT, drug interaction

digoxin: IT, drug interaction

7 ethyl 10 hydroxycamptothecin: IT, drug interaction

indinavir: IT, drug interaction

RN (topotecan) 119413-54-6, 123948-87-8; (protein) 67254-75-5;  
 (paclitaxel) 33069-62-4; (cyclosporin A) 59865-13-3, 63798-73-2;  
 (elacridar) 143664-11-3; (rifampicin) 13292-46-1; (digoxin) 20830-75-5,  
 57285-89-9; (7 ethyl 10 hydroxycamptothecin) 86639-52-3; (indinavir)

150378-17-9, 157810-81-6, 180683-37-8

CN Sn 38; Gf 120918

L154 ANSWER 17 OF 25 ZCAPLUS COPYRIGHT 2004 ACS on STN DUPLICATE 2

AN 2000:824069 ZCAPLUS

DN 134:9341

ED Entered STN: 24 Nov 2000

TI A method of improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and novel pharmaceutical compositions for oral delivery of drugs

IN Schellens, Johannes Henricus Matthias; Schinkel, Alfred Hermanus

PA Het Nederlands Kanker Instituut, Neth.

SO PCT Int. Appl., 25 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K

CC 63-5 (Pharmaceuticals)

FAN.CNT 2

|      | PATENT NO.      | KIND | DATE   | APPLICATION NO. | DATE         |
|------|-----------------|------|--|-----------------|--------------|
| PI   | WO 2000069390   | A2   | 20001123   | WO 2000-NL331   | 20000517 <-- |
|      | WO 2000069390   | A3   | 20011213   |                 |              |
|      | W:              |      | AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM |                 |              |
|      | RW:             |      | GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG   |                 |              |
|      | AU 2000049552   | A5   | 20010205   | AU 2000-49552   | 20000517 <-- |
|      | EP 1189637      | A2   | 20020327   | EP 2000-931720  | 20000517 <-- |
|      | R:              |      | AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO   |                 |              |
|      | US 2002128282   | A1   | 20020912   | US 2001-988285  | 20011119 <-- |
| PRAI | NL 1999-1012066 | A    | 19990517 <--   |                 |              |
|      | NL 1999-1012481 | A    | 19990630 <--   |                 |              |
|      | WO 2000-NL331   | W    | 20000517   |                 |              |

AB A method for increasing the systemic exposure of cells selected from tumor cells and normal cells to an orally administered pharmaceutically active compound, wherein a bioenhancer comprising an inhibitor of breast cancer resistance protein (BCRP)-mediated and/or related drug transport is orally administered concomitantly with said orally administered pharmaceutically active compound, and in which method the inhibitor is administered simultaneously with the pharmaceutical compound Mice were given oral GF120918 (50 mg/kg) 15 min before oral dose of 1 mg/kg topotecan. A profound increased systemic exposure to oral topotecan was observed The increase in the AUC was approx. 6 fold.

ST oral pharmaceutical bioavailability permeation enhancer; breast cancer resistance protein inhibitor pharmaceutical

IT Antitumor agents

(mammary gland; method of improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and novel pharmaceutical compns. for oral delivery of drugs)

IT Antitumor agents

Digestive tract

Drug bioavailability

Permeation enhancers

- (method of improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and novel pharmaceutical compns. for oral delivery of drugs)
- IT Mycotoxins  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(method of improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and novel pharmaceutical compns. for oral delivery of drugs)
- IT Mammary gland  
(neoplasm, inhibitors; method of improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and novel pharmaceutical compns. for oral delivery of drugs)
- IT Drug delivery systems  
(oral; method of improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and novel pharmaceutical compns. for oral delivery of drugs)
- IT 180422-22-4, XR 9051  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(XR 9051; method of improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and novel pharmaceutical compns. for oral delivery of drugs)
- IT 206873-63-4, XR 9576  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(XR 9576; method of improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and novel pharmaceutical compns. for oral delivery of drugs)
- IT 84-65-1D, Anthraquinone, derivs. 253-82-7D, Quinazoline, derivs.  
7689-03-4D, Camptothecin, derivs. 19216-56-9, Prazosin  
65271-80-9, Mitoxantrone 86639-52-3, Sn38 91421-42-0,  
9-Nitrocamptothecin 91421-43-1, 9-Aminocamptothecin  
100286-90-6, Cpt11 118974-02-0, Fumitremorgin c  
123948-87-8, Topotecan. 143664-11-3, GF120918  
149882-10-0, Gg211 169869-90-3, Dx8951f  
203923-89-1, Bnp 1350  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(method of improving bioavailability of orally administered drugs, screening for enhancers of such bioavailability and novel pharmaceutical compns. for oral delivery of drugs)
- L154 ANSWER 18 OF 25 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN DUPLICATE 3
- AN 2000:511112 BIOSIS
- DN PREV200000511112
- TI Role of breast cancer resistance protein in the bioavailability and fetal penetration of **topotecan**.
- AU Jonker, Johan W.; Smit, Johan W.; Brinkhuis, Remco F.; Maliepaard, Marc; Beijnen, Jos H.; Schellens, Jan H. M.; Schinkel, Alfred H. [Reprint author]
- CS Division of Experimental Therapy, Netherlands Cancer Institute, Plesmanlaan 121, 1066 CX, Amsterdam, Netherlands
- SO Journal of the National Cancer Institute (Bethesda), (October 18, 2000) Vol. 92, No. 20, pp. 1651-1656. print.

CODEN: JNCIEQ. ISSN: 0027-8874.

DT Article

LA English

ED Entered STN: 22 Nov 2000

Last Updated on STN: 11 Jan 2002

AB Background and Methods: Breast cancer resistance protein (BCRP/MXR/ABCP) is a multidrug-resistance protein that is a member of the adenosine triphosphate-binding cassette family of drug transporters. BCRP can render tumor cells resistant to the anticancer drugs **topotecan**, mitoxantrone, doxorubicin, and daunorubicin. To investigate the physiologic role of BCRP, we used polarized mammalian cell lines to determine the direction of BCRP drug transport. We also used the BCRP inhibitor GF120918 to assess the role of BCRP in protecting mice against xenobiotic drugs. Bcrp1, the murine homologue of BCRP, was expressed in the polarized mammalian cell lines LLC-PK1 and MDCK-II, and the direction of Bcrp1-mediated transport of **topotecan** and mitoxantrone was determined. To avoid the confounding drug transport provided by P-glycoprotein (P-gp), the roles of Bcrp1 in the bioavailability of **topotecan** and the effect of GF120918 were studied in both wild-type and P-gp-deficient mice and their fetuses. Results: Bcrp1 mediated apically directed transport of drugs in polarized cell lines. When both **topotecan** and GF120918 were administered orally, the bioavailability (i.e., the extent to which a drug becomes available to a target tissue after administration) of **topotecan** in plasma was dramatically increased in P-gp-deficient mice (greater than sixfold) and wild-type mice (greater than ninefold), compared with the control (i.e., vehicle-treated) mice. Furthermore, treatment with GF120918 decreased plasma clearance and hepatobiliary excretion of **topotecan** and increased (re-)uptake by the small intestine. In pregnant GF120918-treated, P-gp-deficient mice, relative fetal penetration of **topotecan** was twofold higher than that in pregnant vehicle-treated mice, suggesting a function for BCRP in the maternal-fetal barrier of the placenta. Conclusions: Bcrp1 mediates apically directed drug transport, appears to reduce drug bioavailability, and protects fetuses against drugs. We propose that strategic application of BCRP inhibitors may thus lead to more effective oral chemotherapy with **topotecan** or other BCRP substrate drugs.

CC Pharmacology - General 22002

Biochemistry studies - General 10060

Pathology - Therapy 12512

Neoplasms - Pathology, clinical aspects and systemic effects 24004

Neoplasms - Therapeutic agents and therapy 24008

Development and Embryology - General and descriptive 25502

IT Major Concepts

Pharmacology; Tumor Biology

IT Chemicals &amp; Biochemicals

**GF-120918: antineoplastic-drug, breast cancer resistance protein inhibitor, drug resistance reversing agent; breast cancer resistance protein: fetal topotecan penetration role, topotecan bioavailability role; topotecan**

**[Hyacinthin]: antineoplastic-drug, bioavailability, fetal penetration, tumor resistance**

ORGN Classifier

Muridae 86375

Super Taxa

Rodentia; Mammalia; Vertebrata; Chordata; Animalia

Organism Name

mouse: animal model, female, fetus

Taxa Notes

Animals, Chordates, Mammals, Nonhuman Vertebrates, Nonhuman Mammals,

Rodents, Vertebrates  
 RN 143664-11-3 (GF-120918)  
 123948-87-8 (topotecan)  
 123948-87-8 (Hyacinthin)  
 38890-82-3 (HYACINTHIN)

L154 ANSWER 19 OF 25 ZCAPLUS COPYRIGHT 2004 ACS on STN

AN 1999:795688 ZCAPLUS

DN 132:35333

ED Entered STN: 17 Dec 1999

TI Multibinding inhibitors of topoisomerase

IN Linsell, Martin S.; Meier-Davis, Susan; Griffin, John H.

PA Advanced Medicine, Inc., USA

SO PCT Int. Appl., 142 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K038-00

ICS A61K039-00; A61K039-44; A61K039-395; A61K051-00; C07K002-00;  
 C07K004-00; G01N033-53; G01N033-543; G01N033-566; C07G011-00

CC 21-2 (General Organic Chemistry)

Section cross-reference(s): 1, 63

FAN.CNT 31

|      | PATENT NO.   | KIND | DATE     | APPLICATION NO.  | DATE         |
|------|--|------|----------|--|--------------|
| PI   | WO 9964054   | A1   | 19991216 | WO 1999-US12908  | 19990608 <-- |
|      | W:   |      |          | AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM |              |
|      | RW:  |      |          | GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG   |              |
|      | US 6288234   | B1   | 20010911 | US 1999-325662   | 19990604 <-- |
|      | CA 2321166   | AA   | 19991216 | CA 1999-2321166  | 19990608 <-- |
|      | AU 9946771   | A1   | 19991230 | AU 1999-46771  | 19990608 <-- |
|      | EP 1085891   | A1   | 20010328 | EP 1999-930179   | 19990608 <-- |
|      | R:   |      |          | AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI   |              |
|      | SG 80631   | A1   | 20010522 | SG 1999-2719   | 19990608 <-- |
|      | SG 90053   | A1   | 20020723 | SG 1999-2944   | 19990608 <-- |
|      | US 6566509   | B1   | 20030520 | US 1999-327899   | 19990608 <-- |
|      | ZA 2000004086  | A    | 20010810 | ZA 2000-4086   | 20000810 <-- |
|      | ZA 2000004558  | A    | 20011130 | ZA 2000-4558   | 20000831 <-- |
|      | ZA 2000004559  | A    | 20020402 | ZA 2000-4559   | 20000831 <-- |
|      | US 2002028943  | A1   | 20020307 | US 2001-760827   | 20010117 <-- |
|      | US 2004023290  | A1   | 20040205 | US 2002-161279   | 20020603 <-- |
|      | US 2003176670  | A1   | 20030918 | US 2002-330381   | 20021227 <-- |
| PRAI | US 1998-88448P   | P    | 19980608 | <--  |              |
|      | US 1998-93072P   | P    | 19980716 | <--  |              |
|      | US 1999-325662   | A3   | 19990604 | <--  |              |
|      | US 1999-327899   | A1   | 19990608 | <--  |              |
|      | US 1999-328071   | B1   | 19990608 | <--  |              |
|      | WO 1999-US12908  | W    | 19990608 | <--  |              |
|      | US 2000-502938   | A1   | 20000211 |  |              |
| AB   | Novel topoisomerase inhibitors that act as multibinding agents, LpXq [where L = a ligand capable of binding to topoisomerase; X = a linker; p = 2-10; q = 1-20; the distance between ligands 2-50 Å], are disclosed. |      |          |  |              |

Combinatorial arrays, methods of synthesis, and methods of assaying the dimeric and multimeric compds. are also embodied by the invention. A number of divalent prophetic examples, derived from substituted fused ring heterocyclic ligands and difunctional linkers, are given. Compds. of this invention are useful in the treatment and prevention of cancer and microbial infections (no data). The multibinding compds. provide greater biol. and/or therapeutic effects than the aggregate of the unlinked ligands due to their multibinding properties (no data). Ligands may include A-62176, A-74932, acridine carboxamides, actinomycin D, AD-312, AD-347, AHMA, AMP-53, amrubicin, amsacrine, anthracyclines, asulacrine, azonafide, azatoxin, BBR-2778, BMJ-43748, BO-2367, bromodeoxyuridine, C-1310, C-1311, CC-131, CJ-12373, CI-937, CI-920 (fostriecin), CP-115953, camptothecin, daunorubicin, doxorubicin, DuP 937 (losoxathrone), DuP 941, elinafide, ellipticine-estradiol (conjugates), elsamitrucin, ER-37328, etoposide, fleroxacin, GI-149893, GL-331, GR-122222X, ICRF-154, ICRF-193, idarubicin, iododoxorubicin, IST-622, KRQ- 10018, intoplicine, lomefloxacin, losoxantrone, m-AMSA, merbarone, meraboin, mitonafide, mitoxantrone, morindone, NCA-0465, NK-109, NK-611, NSC-655649, NSC-665517, NSC-675967, pazelliptine, pazufloxacin, PD-131112, piroxantrone, pyridobenzophenoxazine, S-16020-2, saintopin, sitafloxacin hydrate, SN-22995, sobuzoxane, SR-103, TAS-103, teloxantrone, teniposide, TLC-D-99, top-53, topotecan, tosofloxacin, TRK-710, trovafloxacin, UCE-6, VM-26, VP-16, W5R, WIN-33377, WIN-58161, WIN-645593, WQ-2743, WQ-3034, WR-63320, XR-5942, XR-5000, and 773U82.

- ST dimeric multimeric multibinding topoisomerase inhibitor prepn;  
combinatorial array multibinding topoisomerase inhibitor; anticancer  
multibinding topoisomerase inhibitor prepn; antimicrobial multibinding  
topoisomerase inhibitor prepn
- IT Structure-activity relationship  
(ligand-binding; preparation of multibinding inhibitors of topoisomerase as  
anticancer and antimicrobial agents)
- IT Antimicrobial agents  
Antitumor agents  
Combinatorial library  
Drug delivery systems  
Drug screening  
(preparation of multibinding inhibitors of topoisomerase as anticancer and  
antimicrobial agents)
- IT Anthracyclines  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use);  
BIOL (Biological study); PREP (Preparation); USES (Uses)  
(preparation of multibinding inhibitors of topoisomerase as anticancer and  
antimicrobial agents)
- IT 80449-01-0  
RL: BPR (Biological process); BSU (Biological study, unclassified); MSC  
(Miscellaneous); BIOL (Biological study); PROC (Process)  
(preparation of multibinding inhibitors of topoisomerase as anticancer and  
antimicrobial agents)
- IT 50-28-2DP, Estradiol, dimeric and multimeric derivs. of 50-76-0DP,  
Actinomycin D, dimeric and multimeric derivs. of 59-14-3DP,  
Bromodeoxyuridine, dimeric and multimeric derivs. of 260-94-6DP,  
Acridine, dimeric and multimeric **derivs.** of 478-29-5DP,  
Morindone, dimeric and multimeric derivs. of 519-23-3DP, Ellipticine,  
dimeric and multimeric derivs. of 1506-47-4DP, ICRF-154, dimeric and  
multimeric derivs. of 7689-03-4DP, Camptothecin, dimeric and  
multimeric **derivs.** of 20830-81-3DP, Daunorubicin, dimeric and  
multimeric derivs. of 21416-88-6DP, ICRF-193, dimeric and multimeric  
derivs. of 23214-92-8DP, Doxorubicin, dimeric and multimeric derivs. of  
29767-20-2DP, Teniposide, dimeric and multimeric derivs. of

33419-42-ODP, Etoposide, dimeric and multimeric derivs. of 51264-14-3DP, Amsacrine, dimeric and multimeric derivs. of 54824-17-8DP, Mitonafide, dimeric and multimeric derivs. of 58957-92-9DP, Idarubicin, dimeric and multimeric derivs. of 65222-35-7DP, Pazelliptine, dimeric and multimeric derivs. of 65271-80-9DP, Mitoxantrone, dimeric and multimeric derivs. of 79660-72-3DP, Fleroxacin, dimeric and multimeric derivs. of 83997-75-5DP, Iododoxorubicin, dimeric and multimeric derivs. of 87810-56-8DP, Fostriecin, dimeric and multimeric derivs. of 88303-60-ODP, Losoxantrone, dimeric and multimeric derivs. of 88303-61-1DP, DUP 941, dimeric and multimeric derivs. of 89458-99-1DP, SN 22995, dimeric and multimeric derivs. of 91441-23-5DP, Piroxantrone, dimeric and multimeric derivs. of 91441-48-4DP, CI-937, dimeric and multimeric derivs. of 96404-52-3DP, 773U82, dimeric and multimeric derivs. of 97068-30-9DP, Elsamitrucin, dimeric and multimeric derivs. of 97534-21-9DP, Merbarone, dimeric and multimeric derivs. of 98079-51-7DP, Lomefloxacin, dimeric and multimeric derivs. of 98631-95-9DP, Sobuzoxane, dimeric and multimeric derivs. of 100490-36-6DP, Tosufloxacin, dimeric and multimeric derivs. of 105760-98-3DP, NK-611, dimeric and multimeric derivs. of 110267-81-7DP, Amrubicin, dimeric and multimeric derivs. of 111783-54-1DP, A-62176, dimeric and multimeric derivs. of 122033-48-1DP, WIN-58161, dimeric and multimeric derivs. of 122536-20-3DP, PD 131112, dimeric and multimeric derivs. of 123830-79-5DP, DuP 937, dimeric and multimeric derivs. of **123948-87-8DP**, Topotecan, dimeric and multimeric derivs. of 125974-72-3DP, Intoplicine, dimeric and multimeric derivs. of 127045-41-4DP, Pazufloxacin, dimeric and multimeric derivs. of 127254-12-ODP, Sitaefloxacin, dimeric and multimeric derivs. of 127882-73-9DP, GL-331, dimeric and multimeric derivs. of 128201-92-3DP, IST 622, dimeric and multimeric derivs. of 128420-31-5DP, AD-312, dimeric and multimeric derivs. of 128455-95-8DP, dimeric and multimeric derivs. of 128524-35-6DP, AD 347, dimeric and multimeric derivs. of 129564-92-7DP, Azatoxin, dimeric and multimeric derivs. of 131190-63-1DP, Saintopin, dimeric and multimeric derivs. of 132195-65-4DP, BMY-43748, dimeric and multimeric derivs. of 136440-70-5DP, CP-115953, dimeric and multimeric derivs. of 138154-39-9DP, C-1311, dimeric and multimeric derivs. of 138154-40-2DP, C-1310, dimeric and multimeric derivs. of 140917-67-5DP, Azonafide, dimeric and multimeric derivs. of 143201-31-4DP, NK-109, dimeric and multimeric derivs. of 144675-97-8DP, BBR-2778, dimeric and multimeric derivs. of 145902-76-7DP, dimeric and multimeric derivs. of 146537-07-7DP, WIN-33377, dimeric and multimeric derivs. of 147059-72-1DP, Trovafloxacin, dimeric and multimeric derivs. of 148201-60-9DP, A-74932, dimeric and multimeric derivs. of 148262-19-5DP, Top-53, dimeric and multimeric derivs. of 149809-18-7DP, GI-149893, dimeric and multimeric derivs. of 150829-94-ODP, UCE-6, dimeric and multimeric derivs. of 154310-42-6DP, dimeric and multimeric derivs. of 156340-22-6DP, CJ 12373, dimeric and multimeric derivs. of 162706-37-8DP, Elinafide, dimeric and multimeric derivs. of 172889-50-8DP, NSC 665517, dimeric and multimeric derivs. of 174634-09-4DP, TAS-103, dimeric and multimeric derivs. of 178169-99-8DP, S-16020-2, dimeric and multimeric derivs. of 180981-08-2DP, TRK 710, dimeric and multimeric derivs. of 213185-01-4DP, SR-103, dimeric and multimeric derivs. of

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(target compound; preparation of multibinding inhibitors of topoisomerase as anticancer and antimicrobial agents)

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE



- (1) Brown; Antibiotic and Chemotherapy 7th Ed 1997, P419 ZCAPLUS
- (2) Ehrhardt; Antimicrobial Agents and Chemotherapy 1997, V41(11), P2570 ZCAPLUS
- (3) Fan; J Med Chem 1995, V38(3), P408 ZCAPLUS
- (4) NEORX Corporation; WO 9205802 A1 1992 ZCAPLUS
- (5) Shuker; Science 1996, V274, P1531 ZCAPLUS

L154 ANSWER 20 OF 25 ZCAPLUS COPYRIGHT 2004 ACS on STN

AN 1998:78980 ZCAPLUS

DN 128:254161

ED Entered STN: 11 Feb 1998

TI A general pattern for substrate recognition by P-glycoprotein

AU Seelig, Anna

CS Department of Biophysical Chemistry, Biocenter of the University of Basel, Basel, CH-4056, Switz.

SO European Journal of Biochemistry (1998), 251(1/2), 252-261

CODEN: EJBCAI; ISSN: 0014-2956

PB Springer-Verlag

DT Journal

LA English

CC 6-1 (General Biochemistry)

AB P-glycoprotein actively transports a wide variety of chemical diverse compds. out of the cell. Based on a comparison of a hundred compds. previously tested as P-glycoprotein substrates, we suggest that a set of well-defined structural elements is required for an interaction with P-glycoprotein. The recognition elements are formed by two (type I unit) or three electron donor groups (type II unit) with a fixed spatial separation. Type I units consist of two electron donor groups with a spatial separation of  $2.5 \pm 0.3$  Å. Type II units contain either two electron donor groups with a spatial separation of  $4.6 \pm 0.6$  Å or three electron donor groups with a spatial separation of the outer two groups of  $4.6 \pm 0.6$  Å. All mols. that contain at least one type I or one type II unit are predicted to be P-glycoprotein substrates. The binding to P-glycoprotein increases with the strength and the number of electron donor or hydrogen bonding acceptor groups forming the type I and type II units. Correspondingly, a high percentage of amino acids with hydrogen bonding donor side chains is found in the transmembrane sequences of P-glycoprotein relevant for substrate interaction. Mols. that minimally contain one type II unit are predicted to be inducers of P-glycoprotein over-expression.

ST P glycoprotein substrate inducer structure activity

IT Structure-activity relationship

(P-glycoprotein substrate and inducer; substrate structural patterns for recognition by and induction of over-expression of P-glycoprotein)

IT Hormones, microbial

RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)

(a-factor, substrate; substrate structural patterns for recognition by and induction of over-expression of P-glycoprotein)

IT Hydrogen bond

(acceptor group, in P-glycoprotein; substrate structural patterns for recognition by and induction of over-expression of P-glycoprotein)

IT Biological transport

(by P-glycoprotein; substrate structural patterns for recognition by and induction of over-expression of P-glycoprotein)

IT Conformation

(of electron donor groups in substrates; substrate structural patterns for recognition by and induction of over-expression of P-glycoprotein)

IT Electron donors

(spatial separation in P-glycoprotein; substrate structural patterns for recognition by and induction of over-expression of P-glycoprotein)

- IT P-glycoproteins  
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); PROC (Process)  
 (substrate structural patterns for recognition by and induction of over-expression of P-glycoprotein)
- IT 50-53-3, Chlorpromazine, biological studies 52-86-8, Haloperidol  
 58-40-2, Promazine 59-05-2, Methotrexate 69-05-6, Quinacrine  
 dihydrochloride 390-64-7, Prenylamine 569-61-9, Pararosaniline  
 1622-62-4, Flunitrazepam 5786-21-0, Clozapine 15663-27-1, Cisplatin  
 154531-78-9, CP 117227  
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
 (borderline substrate; substrate structural patterns for recognition by and induction of over-expression of P-glycoprotein)
- IT 50-06-6, Phenobarbital, biological studies 50-55-5, Reserpine 50-76-0, Actinomycin D 53-79-2, Puromycin 57-22-7, Vincristine 64-86-8, Colchicine 114-07-8, Erythromycin 120-58-1, Isosafrole 865-21-4, Vinblastine 13292-46-1, Rifampicin 20830-81-3, Daunorubicin 21829-25-4, Nifedipine 23214-92-8, Doxorubicin 23593-75-1, Clotrimazole 33069-62-4, Taxol 33419-42-0, Etoposide 59467-70-8, Midazolam 152044-53-6, Epothilone A  
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
 (inducer; substrate structural patterns for recognition by and induction of over-expression of P-glycoprotein)
- IT 51-43-4, Epinephrine 72-57-1, Trypan blue 83-60-3, Reserpine acid 4602-84-0, Farnesol 44641-43-2, Cysteine methylester 68000-92-0, Farnesylcysteine 75621-03-3, Chaps  
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
 (non-substrate; substrate structural patterns for recognition by and induction of over-expression of P-glycoprotein)
- IT 7727-37-9, Nitrogen, biological studies 7782-44-7, Oxygen, biological studies  
 RL: BOC (Biological occurrence); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence)  
 (occurrence in substrates; substrate structural patterns for recognition by and induction of over-expression of P-glycoprotein)
- IT 50-02-2, Dexamethasone 50-23-7, Hydrocortisone 50-52-2, Thioridazine 52-39-1, Aldosterone 52-53-9, Verapamil 56-54-2, Quinidine 57-27-2, Morphine, biological studies 57-41-0, Phenytoin 57-83-0, Progesterone, biological studies 58-32-2, Dipyrindamole 58-39-9, Perphenazine 69-23-8, Fluphenazine 85-79-0, Dibucaine 117-89-5, Trifluoperazine 135-67-1, Phenoxazine 146-48-5, Yohimbine 146-54-3, Triflupromazine 481-49-2, Cepharanthine 483-18-1, Emetine 485-71-2, Cinchonidine 749-02-0, Spiperone 1095-90-5, Methadone hydrochloride 1951-25-3, Amiodarone 2001-95-8, Valinomycin 2182-14-1, Vindoline 2468-21-5, Catharanthine 2751-90-8, Tetraphenylphosphonium bromide 2814-89-3, 2901-66-8, Methylreserpate 9002-93-1, Triton X-100 10540-29-1, Tamoxifen 16662-47-8, Gallopamil 17090-79-8, Monensin 19216-56-9, Prazosin 20290-10-2, Morphine 6-glucuronide 20830-75-5, Digoxin 25953-19-9, Cefazolin 42399-41-7, Diltiazem 50679-08-8, Terfenadine 53179-11-6, Loperamide 53772-82-0, cis-Flupenthixol 55985-32-5, Nicardipine 57808-66-9, Domperidone 59865-13-3, Cyclosporin A 62669-70-9, Rhodamine 123 62893-19-0, Cefoperazone 64706-54-3, Bepridil 65271-80-9, Mitoxantrone 69712-56-7, Cefotetan 70288-86-7, Ivermectin 78186-34-2, Bisantrene 90523-31-2, Azidopine 99614-02-5, Ondansetron 104845-40-1, SDB-ethylenediamine 104987-11-3, FK 506

120054-86-6, Dexniguldipine 121584-18-7, SDZ PSC 833 **123948-87-8**  
, Topotecan 130062-64-5 137694-16-7, BIBW22BS 140945-01-3, S 9788  
142716-85-6, CP 100356 **143664-11-3**, GF 120918

RL: BPR (Biological process); BSU (Biological study, unclassified); PRP  
(Properties); BIOL (Biological study); PROC (Process)

(substrate; substrate structural patterns for recognition by and  
induction of over-expression of P-glycoprotein)

RE.CNT 68 THERE ARE 68 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Akiyama, S; Mol Pharmacol 1988, V33, P144 ZCAPLUS
- (2) Azzaria, M; Mol Cell Biol 1989, V9, P5289 ZCAPLUS
- (3) Beck, W; Biochem Biophys Res Commun 1988, V153, P959 ZCAPLUS
- (4) Bhat, U; Mol Pharmacol 1995, V48, P682 ZCAPLUS
- (5) Board, P; FEBS Lett 1993, V315, P298 ZCAPLUS
- (6) Bollag, D; Cancer Res 1995, V55, P2325 ZCAPLUS
- (7) Borst, P; Annu Rev Microbiol 1995, V49, P427 ZCAPLUS
- (8) Buckingham, L; Int J Cancer 1996, V65, P74 ZCAPLUS
- (9) Chen, C; Cell 1986, V47, P381 ZCAPLUS
- (10) Chin, K; J Biol Chem 1990, V265, P221 ZCAPLUS
- (11) Cordon-Cardo, C; J Histochem Cytochem 1990, V38, P1277 ZCAPLUS
- (12) Cordon-Cardo, C; Proc Natl Acad Sci USA 1990, V86, P695
- (13) Descoteaux, S; Gene (Amst) 1995, V164, P179 ZCAPLUS
- (14) Doige, C; Annu Rev Microbiol 1993, V47, P291 ZCAPLUS
- (15) Ferry, D; Eur J Cancer 1996, V32A, P1070 ZCAPLUS
- (16) Ford, J; Eur J Cancer 1996, V32A, P991 ZCAPLUS
- (17) Ford, J; Mol Pharmacol 1988, V35, P105
- (18) Genne, P; Cancer Res 1992, V52, P2797 ZCAPLUS
- (19) Germann, U; Eur J Cancer 1996, V32A, P927 ZCAPLUS
- (20) Gosland, M; Cancer Res 1989, V49, P6901 ZCAPLUS
- (21) Gosland, M; Cancer Res 1993, V53, P5382 ZCAPLUS
- (22) Gottesman, M; Annu Rev Biochem 1993, V62, P385 ZCAPLUS
- (23) Gupta, S; J Clin Immunol 1993, V13, P289 ZCAPLUS
- (24) Hait, W; Biochem Pharmacol 1993, V45, P401 ZCAPLUS
- (25) Higgins, C; Annu Rev Cell Biol 1992, V8, P67 ZCAPLUS
- (26) Hsu, S; Mol Cell Biol 1990, V10, P3596 ZCAPLUS
- (27) Huwiler, J; Br J Pharmacol 1996, V118, P1879
- (28) Hyafil, F; Cancer Res 1993, V53, P4595 ZCAPLUS
- (29) Ichikawa, M; J Biol Chem 1991, V266, P903 ZCAPLUS
- (30) Kajiji, S; Biochemistry 1994, V33, P5041 ZCAPLUS
- (31) Kast, C; Biochemistry 1995, V34, P4402 ZCAPLUS
- (32) Liminga, G; Exp Cell Res 1994, V212, P291 ZCAPLUS
- (33) Liu, Z; Mol Pharmacol 1996, V50, P482 ZCAPLUS
- (34) Loe, D; Biochim Biophys Acta 1994, V1190, P72 ZCAPLUS
- (35) Marks, D; Br J Haematol 1996, V95, P587 ZCAPLUS
- (36) McClean, S; Biochim Biophys Acta 1993, V1177, P117 ZCAPLUS
- (37) Mukhopadhyay, T; J Natl Cancer Inst 1988, V80, P269 ZCAPLUS
- (38) Muller, C; Biochem Pharmacol 1992, V43, P2091 ZCAPLUS
- (39) Nare, B; Mol Pharmacol 1994, V45, P1145 ZCAPLUS
- (40) Pearce, H; Proc Natl Acad Sci USA 1989, V86, P5128 ZCAPLUS
- (41) Pickarz, R; J Biol Chem 1993, V268, P7613
- (42) Qian, X; J Biol Chem 1990, V265, P18753 ZCAPLUS
- (43) Rao, U; Biochem Pharmacol 1994, V48, P287 ZCAPLUS
- (44) Raymond, M; Mol Cell Biol 1994, V14, P277 ZCAPLUS
- (45) Saeki, T; J Biol Chem 1993, V268, P6077 ZCAPLUS
- (46) Safa, A; Biochem Biophys Res Commun 1994, V202, P606 ZCAPLUS
- (47) Safa, A; Biochemistry 1994, V33, P256 ZCAPLUS
- (48) Safa, A; Cancer Invest 1993, V11, P46 ZCAPLUS
- (49) Schinkel, A; J Clin Invest 1996, V97, P2517 ZCAPLUS
- (50) Schuetz, E; Mol Pharmacol 1995, V49, P311
- (51) Seelig, A; Proc Natl Acad Sci USA 1994, V91, P68 ZCAPLUS

- (52) Shapiro, A; J Biol Chem 1995, V270, P16167 MEDLINE
- (53) Sharom, F; J Biol Chem 1993, V268, P24197 ZCAPLUS
- (54) Tamai, I; J Biol Chem 1991, V266, P16796 ZCAPLUS
- (55) Thiebaut, F; J Histochem Cytochem 1989, V37, P159 ZCAPLUS
- (56) Thiebaut, F; Proc Natl Acad Sci USA 1987, V84, P7735 ZCAPLUS
- (57) Thimmaiah, K; Cancer Commun 1990, V2, P249 ZCAPLUS
- (58) Uchiumi, T; Cell Growth Differ 1993, V4, P147 ZCAPLUS
- (59) Ueda, K; J Biol Chem 1992, V267, P24248 ZCAPLUS
- (60) Urbatsch, I; J Biol Chem 1995, V270, P19383 ZCAPLUS
- (61) Vinogradov, S; Hydrogen bonding 1971, P11
- (62) Yusa, K; Cancer Res 1989, V49, P5002 ZCAPLUS
- (63) Zamora, J; Mol Pharmacol 1988, V33, P454 ZCAPLUS
- (64) Zhang, L; J Biol Chem 1994, V269, P15973 ZCAPLUS
- (65) Zhang, L; J Biol Chem 1995, V39, P22859
- (66) Zhang, X; J Biol Chem 1995, V270, P5441 ZCAPLUS
- (67) Zhang, X; Oncol Res 1994, V6, P291 ZCAPLUS
- (68) Zordan-Nudo, T; Cancer Res 1993, V53, P5994 ZCAPLUS

L154 ANSWER 21 OF 25 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN

AN 1997:517625 BIOSIS

DN PREV199799816828

TI Cellular responses to methyl-N-(4-(9-acridinylamino)-2-methoxyphenyl)carbamate hydrochloride, an analogue of amsacrine active against non-proliferating cells.

AU Moreland, N.; Finlay, G. J.; Dragunow, M.; Holdaway, K. M.; Baguley, B. C.  
[Reprint author]

CS Cancer Research Lab., Univ. Auckland Sch. Med., Private Bag 92019,  
Auckland 1000, New Zealand

SO European Journal of Cancer, (1997) Vol. 33, No. 10, pp. 1668-1676.  
CODEN: EJCAEL. ISSN: 0959-8049.

DT Article

LA English

ED Entered STN: 10 Dec 1997

Last Updated on STN: 10 Dec 1997

AB The **acridine derivative** m-AMCA (methyl-N-(4-(9-acridinylamino)-2-methoxyphenyl)carbamate hydrochloride), a carbamate analogue of the topoisomerase II poison amsacrine, is distinguished by its high cytotoxicity against non-cycling tumour cells. We compared the response of cultured Lewis lung carcinoma cells to b-AMCA, amsacrine and the topoisomerase I poison camptothecin. The DNA polymerase inhibitor aphidicolin reversed the cytotoxicity of camptothecin fully, that of amsacrine partially, and that of m-AMCA minimally. The ability of m-AMCA to induce the enzyme poly(ADP-ribose)polymerase (PARP) was markedly lower than that of camptothecin or amsacrine. Cell cycle responses to m-AMCA and amsacrine were similar, with slowing of progress through S-phase and arrest in G-2-phase. These cell cycle changes were also observed when plateau phase cultures were exposed to drug for 1 h, washed free of drug and cultured in fresh medium, with m-AMCA having a more pronounced effect than amsacrine and camptothecin having no effect. We also examined the role of p53 protein in the response using cultured human H460 cells. Both mAMCA and amsacrine induced p53 protein expression in proliferating but not in non-proliferating H460 cells, and induced p21-WAF1 regardless of proliferation status. Both induced G-1-phase cell cycle arrest. It is suggested that two cytotoxicity mechanisms can be distinguished using these drugs. The first is specific for S-phase cells, is reversed by aphidicolin and induces PARP activity. The second is cell cycle non-specific, does not induce PARP and is unaffected by aphidicolin. Camptothecin activates only the first, m-AMCA primarily the second and amsacrine activates both.

CC Cytology - Animal 02506

Cytology - Human 02508  
Biochemistry studies - General 10060  
Biochemistry studies - Proteins, peptides and amino acids 10064  
Enzymes - Physiological studies 10808  
Pathology - Therapy 12512  
Respiratory system - Pathology 16006  
Pharmacology - Drug metabolism and metabolic stimulators 22003  
Pharmacology - Respiratory system 22030  
Neoplasms - Neoplastic cell lines 24005  
Neoplasms - Biochemistry 24006  
Neoplasms - Therapeutic agents and therapy 24008

IT Major Concepts  
Cell Biology; Enzymology (Biochemistry and Molecular Biophysics);  
Oncology (Human Medicine, Medical Sciences); Pharmacology; Pulmonary  
Medicine (Human Medicine, Medical Sciences)

IT Chemicals & Biochemicals  
AMSACRINE; TOPOISOMERASE II; **CAMPTOTHECIN**; POLY-(ADP-RIBOSE)  
POLYMERASE

IT Miscellaneous Descriptors  
AMSACRINE; AMSACRINE ANALOGUE; ANTINEOPLASTIC-DRUG; CAMPTOTHECIN; DRUG  
TREATMENT; DRUG-INDUCED ACTIVATION; H-460 CELL LINE; HUMAN LUNG CANCER  
CELL LINE; IN-VITRO MODEL SYSTEM; LLTC CELL LINE; METHYL-N-(4-(9-  
ACRIDINYLAMINO)-2-METHOXYPHENYL)CARBAMATE HYDROCHLORIDE; MOUSE LEWIS  
LUNG CARCINOMA CELL LINE; PHARMACOLOGY; POLY-(ADP-RIBOSE) POLYMERASE;  
TOPOISOMERASE I POISON; TOPOISOMERASE II POISON; TUMOR BIOLOGY

ORGN Classifier  
Hominidae 86215  
Super Taxa  
Primates; Mammalia; Vertebrata; Chordata; Animalia  
Organism Name  
Hominidae  
Taxa Notes  
Animals, Chordates, Humans, Mammals, Primates, Vertebrates

ORGN Classifier  
Muridae 86375  
Super Taxa  
Rodentia; Mammalia; Vertebrata; Chordata; Animalia  
Organism Name  
Muridae  
Taxa Notes  
Animals, Chordates, Mammals, Nonhuman Vertebrates, Nonhuman Mammals,  
Rodents, Vertebrates

RN 51264-14-3 (AMSACRINE)  
142805-56-9 (TOPOISOMERASE II)  
7689-03-4 (CAMPTOTHECIN)  
9055-67-8 (POLY-(ADP-RIBOSE) POLYMERASE)

L154 ANSWER 22 OF 25 ZCAPLUS COPYRIGHT 2004 ACS on STN  
AN 1996:741460 ZCAPLUS  
DN 126:84167  
ED Entered STN: 18 Dec 1996  
TI Selective induction of apoptosis in Hep 3B cells by topoisomerase I  
inhibitors: evidence for protease-dependent pathway that does not activate  
cysteine protease P32  
AU Adjei, Philip N.; Kaufmann, Scott H.; Leung, Wai-Yee; Mao, Fei; Gores,  
Gregory J.  
CS Div. Gastrointestinal Int. Med., Mayo Clinic Found., Rochester, MN, 55905,  
USA  
SO Journal of Clinical Investigation (1996), 98(11), 2588-2596  
CODEN: JCINAO; ISSN: 0021-9738

PB Rockefeller University Press

DT Journal

LA English

CC 1-6 (Pharmacology)

AB Progress in the treatment of hepatocellular carcinoma (HCC), a common tumor worldwide, has been disappointing. Inhibitors of topoisomerases are being widely studied as potential inducers of tumor cell apoptosis. The authors aims were to determine whether topoisomerase-directed drugs would induce apoptosis in a human HCC cell line (Hep 3B) and, if so, to investigate the mechanism. The topoisomerase I poison camptothecin (CPT) induced apoptosis of Hep 3B cells in a time- and concentration-dependent manner.

In contrast, the topoisomerase II poison etoposide failed to induce apoptosis despite the apparent stabilization of topoisomerase II-DNA complexes. Unexpectedly, CPT-induced apoptosis in this cell type occurred without any detectable cleavage of poly(ADP-ribose) polymerase or lamin B, polypeptides that are commonly cleaved in other cell types undergoing apoptosis. Likewise, Hep 3B cell apoptosis occurred without a detectable increase in interleukin-1 $\beta$ -converting enzyme (ICE)-like or cysteine protease P32 (CPP32)-like protease activity. In contrast, trypsin-like protease activity (cleavage of Boc-Val-Leu-Lys-chloromethylaminocoumarin in situ) increased threefold in cells treated with CPT but not etoposide. Tosyl-lysyl chloromethyl ketone inhibited the trypsin-like protease activity and diminished CPT-induced apoptosis. These data demonstrate that (a) apoptosis is induced in Hep 3B cells after stabilization of topoisomerase I-DNA complexes but not after stabilization of topoisomerase II-DNA complexes as measured by alkaline filter elution; (b) Hep 3B cell apoptosis occurs without activation of ICE-like and CPP32-like protease activity; and (c) a trypsin-like protease activity appears to contribute to apoptosis in this cell type.

ST apoptosis hepatocellular carcinoma topoisomerase inhibitor protease

IT Liver, neoplasm

(hepatoma, inhibitors; selective induction of apoptosis in human hepatocellular carcinoma Hep 3B cells by topoisomerase I inhibitors and evidence for protease-dependent pathway that does not activate cysteine protease P32)

IT Antitumor agents

(hepatoma; selective induction of apoptosis in human hepatocellular carcinoma Hep 3B cells by topoisomerase I inhibitors and evidence for protease-dependent pathway that does not activate cysteine protease P32)

IT Apoptosis

(selective induction of apoptosis in human hepatocellular carcinoma Hep 3B cells by topoisomerase I inhibitors and evidence for protease-dependent pathway that does not activate cysteine protease P32)

IT 143180-75-0, DNA topoisomerase I

RL: ADV (Adverse effect, including toxicity); BIOL (Biological study) (inhibitors; selective induction of apoptosis in human hepatocellular carcinoma Hep 3B cells by topoisomerase I inhibitors and evidence for protease-dependent pathway that does not activate cysteine protease P32)

IT **260-94-6D**, Acridine, pyrazolo-fused **derivative** 7689-03-4,

Camptothecin 33419-42-0, Etoposide 86639-52-3, SN-38

**123948-87-8**, Topotecan

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(selective induction of apoptosis in human hepatocellular carcinoma Hep 3B cells by topoisomerase I inhibitors and evidence for

protease-dependent pathway that does not activate cysteine protease  
P32)

- IT 122191-40-6, Interleukin-1 $\beta$ -converting enzyme 169592-57-8, CPP32  
proteinase  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL  
(Biological study); PROC (Process)  
(selective induction of apoptosis in human hepatocellular carcinoma Hep  
3B cells by topoisomerase I inhibitors and evidence for  
protease-dependent pathway that does not activate cysteine protease  
P32)
- IT 37259-58-8, Serine proteinase  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL  
(Biological study); PROC (Process)  
(trypsin-like; selective induction of apoptosis in human hepatocellular  
carcinoma Hep 3B cells by topoisomerase I inhibitors and evidence for  
protease-dependent pathway that does not activate cysteine protease  
P32)

L154 ANSWER 23 OF 25 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN  
AN 1996:113207 BIOSIS  
DN PREV199698685342

TI Interaction of cytostatics and chemosensitizers with the dextriguldipine  
binding site on P-glycoprotein.

AU Boer, Rainer [Reprint author]; Ulrich, Wolf-Ruediger; Haas, Sabine;  
Borchers, Christoph; Gekeler, Volker; Boss, Hildegaard; Przybylski,  
Michael; Schoedl, Angelika

CS Byk Gulden, Byk Gulden-Str. 2, D-78467 Konstanz, Germany

SO European Journal of Pharmacology, (1996) Vol. 295, No. 2-3, pp. 253-260.  
CODEN: EJPHAZ. ISSN: 0014-2999.

DT Article

LA English

ED Entered STN: 12 Mar 1996

Last Updated on STN: 13 Mar 1996

AB The interaction of cytostatics and chemosensitizers with the  
dextriguldipine binding site of P-glycoprotein was investigated in  
photoaffinity labeling experiments. A tritiated azidoderivative of the  
chemosensitizer dextriguldipine with dihydropyridine structure,  
(3H)B9209-005, was used to irreversibly label P-glycoprotein. The  
apparent affinity of cytostatics and chemosensitizers to this binding site  
was estimated from labeling experiments in the presence of increasing  
concentrations of compounds. From the cytostatics tested, the vinca  
alkaloids and taxol showed the highest affinity, anthracyclins possessed  
moderate affinity while methotrexate, ara C and camptothecin, cytostatics  
not involved in P-glycoprotein-mediated multidrug resistance, were almost  
inactive. The chemosensitizers **GF 120918**, cyclosporin  
A and SDZ PSC-833 inhibited photoincorporation with the highest potency.  
Steep dose-inhibition curves were obtained with the cyclic peptides and  
S9788, indicating that these compounds may bind allosterically to a  
separate binding site. Compounds with dihydropyridine structure with or  
without chemosensitizing potency were also tested and some  
structure-activity relationships could be derived from the data. Our data  
show that inhibition of photoaffinity labeling by (3H)B9209-005 is a  
valuable and reliable system for measuring the interaction with and  
potency of chemosensitizing compounds at P-glycoprotein. Furthermore,  
data obtained in this test system are well suited to investigate  
structure-activity relationships for chemosensitizers at P-glycoprotein.  
In addition cytostatics underlying P-glycoprotein-mediated multidrug  
resistance can be identified.

CC Cytology - Human 02508

Biochemistry studies - General 10060

Biochemistry studies - Proteins, peptides and amino acids 10064  
 Biochemistry studies - Carbohydrates 10068  
 Biophysics - Membrane phenomena 10508  
 Blood - Lymphatic tissue and reticuloendothelial system 15008  
 Pharmacology - Muscle system 22022  
 Neoplasms - Therapeutic agents and therapy 24008  
 IT Major Concepts  
     Blood and Lymphatics (Transport and Circulation); Cell Biology;  
     Membranes (Cell Biology); Oncology (Human Medicine, Medical Sciences);  
     Pharmacology  
 IT Chemicals & Biochemicals  
     METHOTREXATE; CYTARABINE; **CAMPTOTHECIN**; CYCLOSPORINE A; SDZ  
     PSC-833  
 IT Miscellaneous Descriptors  
     **CAMPTOTHECIN**; CYCLOSPORINE A; CYTARABINE; **GF 120918**  
     ; HUMAN LYMPHOBLASTOID CELLS; METHOTREXATE; MULTIDRUG RESISTANCE; SDZ  
     PSC-833  
 ORGN Classifier  
     Hominidae 86215  
     Super Taxa  
         Primates; Mammalia; Vertebrata; Chordata; Animalia  
     Organism Name  
         Hominidae  
     Taxa Notes  
         Animals, Chordates, Humans, Mammals, Primates, Vertebrates  
 RN 59-05-2 (METHOTREXATE)  
     147-94-4 (CYTARABINE)  
     7689-03-4 (CAMPTOTHECIN)  
     59865-13-3 (CYCLOSPORINE A)  
     121584-18-7 (SDZ PSC-833)  
 L154 ANSWER 24 OF 25 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN  
 AN 1995:161503 BIOSIS  
 DN PREV199598175803  
 TI DNA topoisomerases: Genome gate-keepers and their intruders, anticancer  
     and antibacterial drugs.  
 AU Pommier, Yves  
 CS Lab. Mol. Pharmacol., Build. 37, Room 5C25, Natl. Cancer Inst., Natl.  
     Inst. Health, Bethesda, MD 20892, USA  
 SO M-S (Medecine Sciences), (1994) Vol. 10, No. 10, pp. 953-955.  
     ISSN: 0767-0974.  
 DT Article  
     Editorial  
 LA French  
 ED Entered STN: 11 Apr 1995  
     Last Updated on STN: 12 Apr 1995  
 CC Genetics - Plant 03504  
     Genetics - Animal 03506  
     Biochemistry studies - General 10060  
     Biochemistry studies - Nucleic acids, purines and pyrimidines 10062  
     Biochemistry studies - Proteins, peptides and amino acids 10064  
     Enzymes - Physiological studies 10808  
     Pharmacology - Drug metabolism and metabolic stimulators 22003  
     Neoplasms - Biochemistry 24006  
     Neoplasms - Therapeutic agents and therapy 24008  
     Genetics of bacteria and viruses 31500  
     Medical and clinical microbiology - Bacteriology 36002  
     Chemotherapy - Antibacterial agents 38504  
     Invertebrata: comparative, experimental morphology, physiology and  
     pathology - Insecta: physiology 64076



IT Major Concepts  
Biochemistry and Molecular Biophysics; Enzymology (Biochemistry and  
Molecular Biophysics); Genetics; Pharmacology; Physiology; Tumor  
Biology

IT Chemicals & Biochemicals  
DNA TOPOISOMERASES; VEPESID; DOXORUBICIN; MITOXANTRONE;  
**ACRIDINE**; ELLIPTICINE; **CAMPTOTHECIN**; TAXOL; TAXOTERE;  
CISPLATIN

IT Miscellaneous Descriptors  
ACRIDINE; ANTINEOPLASTIC-DRUG; CAMPTOTHECIN; CISPLATIN; DNA  
RECOMBINATION; DOXORUBICIN; ELLIPTICINE; ENZYME INHIBITOR-DRUG; ENZYME  
MUTATION; EUKARYOTIC CELLS; MITOXANTRONE; QUINOLONES; TAXOL; TAXOTERE;  
VEPESID

ORGN Classifier  
Diptera 75314  
Super Taxa  
Insecta; Arthropoda; Invertebrata; Animalia  
Organism Name  
Drosophila  
Taxa Notes  
Animals, Arthropods, Insects, Invertebrates

ORGN Classifier  
Fungi 15000  
Super Taxa  
Plantae  
Organism Name  
fungi  
yeast  
Taxa Notes  
Fungi, Microorganisms, Nonvascular Plants, Plants

ORGN Classifier  
Mammalia 85700  
Super Taxa  
Vertebrata; Chordata; Animalia  
Organism Name  
mammals  
Mammalia  
Taxa Notes  
Animals, Chordates, Mammals, Nonhuman Vertebrates, Nonhuman Mammals,  
Vertebrates

ORGN Classifier  
Poxviridae 03110  
Super Taxa  
dsDNA Viruses; Viruses; Microorganisms  
Organism Name  
vaccinia virus  
Taxa Notes  
Double-Stranded DNA Viruses, Microorganisms, Viruses

ORGN Classifier  
Retroviridae 03305  
Super Taxa  
DNA and RNA Reverse Transcribing Viruses; Viruses; Microorganisms  
Organism Name  
human immunodeficiency virus  
Taxa Notes  
DNA and RNA Reverse Transcribing Viruses, Microorganisms, Viruses

RN 80449-01-0D (DNA TOPOISOMERASES)  
33419-42-0 (VEPESID)  
23214-92-8 (DOXORUBICIN)  
65271-80-9 (MITOXANTRONE)

260-94-6 (ACRIDINE)  
519-23-3 (ELLIPTICINE)  
7689-03-4 (CAMPTOTHECIN)  
33069-62-4 (TAXOL)  
114977-28-5 (TAXOTERE)  
15663-27-1 (CISPLATIN)

L154 ANSWER 25 OF 25 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN  
AN 1995:19629 BIOSIS  
DN PREV199598033929  
TI DNA topoisomerases and topoisomerase inhibitors.  
AU Giaccone, G.  
CS Dep. Oncol., Free Univ. Hosp., De Boelelaan 1117, 1081 HV Amsterdam,  
Netherlands  
SO Pathologie Biologie, (1994) Vol. 42, No. 4, pp. 346-352.  
CODEN: PABIAQ. ISSN: 0369-8114.  
DT Article  
General Review; (Literature Review)  
LA English  
ED Entered STN: 11 Jan 1995  
Last Updated on STN: 23 Feb 1995  
AB DNA topoisomerases are ubiquitous nuclear enzymes, essential for several  
steps of DNA metabolism. They have recently been shown to be specific  
targets of a number of anticancer agents. In this review are discussed  
the most recent discoveries in the physiology and the molecular biology of  
DNA topoisomerases, and the mechanism of interaction with drugs. In  
addition, alterations of DNA topoisomerases are also described, as  
potential responsible of drug resistance.  
CC Cytology - Animal 02506  
Cytology - Human 02508  
Genetics - Animal 03506  
Genetics - Human 03508  
Biochemistry studies - General 10060  
Biochemistry studies - Nucleic acids, purines and pyrimidines 10062  
Biochemistry studies - Proteins, peptides and amino acids 10064  
Enzymes - Physiological studies 10808  
Pathology - Necrosis 12510  
Pathology - Therapy 12512  
Metabolism - Nucleic acids, purines and pyrimidines 13014  
Pharmacology - Drug metabolism and metabolic stimulators 22003  
Pharmacology - Clinical pharmacology 22005  
Neoplasms - Pathology, clinical aspects and systemic effects 24004  
Neoplasms - Therapeutic agents and therapy 24008  
IT Major Concepts  
Biochemistry and Molecular Biophysics; Cell Biology; Enzymology  
(Biochemistry and Molecular Biophysics); Genetics; Metabolism; Oncology  
(Human Medicine, Medical Sciences); Pharmacology  
IT Chemicals & Biochemicals  
DNA TOPOISOMERASES; TOPOISOMERASE; DOXORUBICIN; DAUNORUBICIN;  
IDARUBICIN; VP-16; VM-26; AMSACRINE; **ACRIDINE**; MITOXANTRONE;  
BISANTRENE; MERBARONE; SURAMIN; ACTINOMYCIN D; **TOPOTECAN**;  
9-AMINO-CAMPTOTHECIN  
IT Miscellaneous Descriptors  
ACRIDINE CARBOXAMIDE; ACTINOMYCIN D; AMSACRINE; ANTHRAPHYRAZOLE;  
ANTINEOPLASTIC-DRUG; ANTINEOPLASTIC-DRUG VM-26; BISANTRENE; CANCER;  
CELL DEATH; DAUNORUBICIN; DNA METABOLISM; DOXORUBICIN; IDARUBICIN;  
MERBARONE; MITOXANTRONE; SURAMIN; **TOPOTECAN**; VP-16;  
2-METHYL-9-HYDROXYELLIPTICINE; 9-AMINO-CAMPTOTHECIN  
ORGN Classifier  
Hominidae 86215

Super Taxa  
     Primates; Mammalia; Vertebrata; Chordata; Animalia  
 Organism Name  
     human  
 Taxa Notes  
     Animals, Chordates, Humans, Mammals, Primates, Vertebrates  
 ORGN Classifier  
     Rodentia     86265  
 Super Taxa  
     Mammalia; Vertebrata; Chordata; Animalia  
 Organism Name  
     rodent  
     Rodentia  
 Taxa Notes  
     Animals, Chordates, Mammals, Nonhuman Vertebrates, Nonhuman Mammals,  
     Rodents, Vertebrates  
 RN   80449-01-0D (DNA TOPOISOMERASES)  
       80449-01-0 (TOPOISOMERASE)  
       23214-92-8 (DOXORUBICIN)  
       20830-81-3 (DAUNORUBICIN)  
       58957-92-9 (IDARUBICIN)  
       33419-42-0 (VP-16)  
       29767-20-2 (VM-26)  
       51264-14-3 (AMSACRINE)  
       260-94-6 (ACRIDINE)  
       65271-80-9 (MITOXANTRONE)  
       78186-34-2 (BISANTRENE)  
       97534-21-9 (MERBARONE)  
       145-63-1 (SURAMIN)  
       50-76-0 (ACTINOMYCIN D)  
       123948-87-8 (TOPOTECAN)  
       91421-43-1 (9-AMINO-CAMPTOTHECIN)

=> b home

FILE 'HOME' ENTERED AT 11:06:11 ON 15 JUN 2004